

JSMDC

Journal of Sharif Medical & Dental College



02/ Vol. 11
December
2025



EDITORIAL BOARD

Editor-in-Chief

Muhammad Adnan Khan

Editor

Maria Aslam

Associate Editors

Muhammad Shahid Iqbal

Saima Inam

Salman Bakhtiar

Uzma Ahsan

Assistant Editors

Amna Iqbal Butt

Uzma Shahid

Members National

A. Majeed Chaudhry

Abid Ashar

Arif Tajammul

Asaf Beg Mirza

Javed Akram

Nausheen Raza

Nazia Yazdanie

Rakhshanda Tayyab

Members International

Adnan H. Siddiqui (USA)

Amina Mirza (UK)

Rizwan Hamid (UK)

Theodore Tyberg (USA)

Ulrich Sigwart (SWL)

Bibliographer

Laila Afzal

Statistician

Muhammad Ibrahim

Editorial Assistant

Khadija Haroon

JSMDC

Journal of Sharif Medical & Dental College

Volume 11

Number 02

December 2025

EDITORIAL

Rising Rates of Antimicrobial Resistance: A Slow Crisis Demanding Urgent Action
Maria Aslam

60-61

ORIGINAL ARTICLES

Association of HbA1c Levels with Complexity of Coronary Artery Disease in Diabetic Patients Presenting with Acute Coronary Syndrome

62-67

Umer Shafiq, Ismail Ahmed Khan, Sohail Aziz, Kashif ur Rehman, Tayyeb Muhammad, Farman Ullah

Efficacy of Lacosamide in Patients with Diabetic Neuropathy

68-72

Awais Akram, Shahzad Alam Khan, Umair Ahmad

Intestinal Parasitic Infections among Children and Young Adolescents Visiting Provincial Headquarter Hospital in Gilgit, Pakistan and their Associated Factors

73-78

Uzma Ali, Sadaf Nasir, Adnan Qureshi, Aqsa Aslam, Namra Yunus, Zakia Wahid

Diagnostic Accuracy of Retrograde Urethrogram in Determination of Urethral Stricture taking Cystoscopy as Gold Standard

79-83

Rafia Irum, Aneesa Qayyum, Hira Bushra, Hamna Mazhar

Comparison of Effectiveness between Diclofenac Suppositories and Injection Diclofenac in Abdominal Surgeries

84-89

Muhammad Abu Usama, Fizza Laraib, Farhana Manzoor, Maham Fatima, Syed Suleman Zafar Jafary, Hafiz Zain-ul-Abdeen

Association of Radial Artery Diameter with Radial Artery Spasm in Patients Undergoing Transradial Cardiac Catheterization

90-95

Ismail Ahmed Khan, Umer Shafiq, Sohail Aziz, Zohair Aziz, Wajahat Lodhi, Waleed Hassan, Asif Manzoor

Early Clinical Exposure through Modular Curriculum in Undergraduate Medical Students: Student Experiences

96-101

Lamia Yusuf, Aqsa Aslam, Maaz Ahmed, Muhammad Fayyaz Atif, Muhammad Hashim Khan, Muhammad Bin Ahsan

Ticagrelor versus Clopidogrel in Complex Percutaneous Coronary Intervention in terms of Mortality and Stent Thrombosis

102-106

Mujaddid Mudassir, Tanvir Ahmad Raja, Hamza Azhar Ghauri, Muhammad Umar Farooq, Ephraim Sheraz, Muhammad Husnain Yousaf

JSMDC is recognized by:

- Higher Education Commission (HEC), Pakistan
- Pakistan Medical & Dental Council (PMDC)

JSMDC is indexed in:

- Directory of Open Access Journals (DOAJ)
- Asian Science Citation Index (ASCI)
- Index Medicus for the Eastern Mediterranean Region (IMEMR)
- PakMediNet

Editorial Correspondence should be addressed to

Editor JSMDC
Sharif Medical & Dental College
Sharif Medical City Road,
Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Tel

(+92-42) 111-123-786

Fax

(+92-42)37860122

E-mail

editors@jsmdc.pk

Website

www.jsmdc.pk

Annual Subscription Fee:

PKR. 2000/-

INDICES**III**

Content Index	107-108
Subject Index	109-110
Author Index	111

INSTRUCTIONS TO AUTHORS**IV**

Rising Rates of Antimicrobial Resistance: A Slow Crisis Demanding Urgent Action

Maria Aslam

Antimicrobial resistance (AMR) is no longer a distant threat; it is a crisis that is encountered in our hospitals and communities. Across the world, patients present with infections that no longer respond to previously effective antibiotics. This silent but relentless threat is undermining decades of medical progress and putting even routine procedures at risk. According to the World Health Organization (WHO), resistance to commonly used antibiotics has become widespread, with alarming levels seen in bacteria such as *Escherichia coli*, *Klebsiella pneumoniae*, and *Staphylococcus aureus*.¹ The WHO warns that the situation is deteriorating faster than expected, particularly in low- and middle-income countries where antibiotic misuse and poor infection control practices are common. The Global Burden of Disease (GBD) 2021 and Antimicrobial Resistance Collaborators, led by Naghavi and colleagues, analyzed data from 204 countries and concluded that antimicrobial resistance was directly responsible for 1.27 million deaths and associated with nearly 5 million deaths in 2019. The study projected a continued rise by 2050 if urgent global action is not taken.²

Antimicrobial resistance is not just a medical problem, it is a social and economic challenge. Antimicrobial resistance ranks as the third most common cause of mortality in Pakistan. Around 59,200 deaths are directly and an additional 221,300 deaths are indirectly caused by AMR in Pakistan. In Pakistan, drug resistance is most commonly seen in *Mycobacterium tuberculosis*, *Staphylococcus aureus*, *Salmonella enterica*, *Enterobacterales*, and non-*Enterobacterales*.³

The etiology of this escalating issue is multifactorial. The major factor is the over-the-counter availability and inappropriate prescription of antibiotics. Patients often demand antibiotics for viral infections, while

physicians may prescribe them empirically without diagnostic evidence of bacterial etiology. In the agricultural sector, antibiotics are still used for growth promotion and prophylaxis in livestock, which contributes to resistant bacterial strains entering the human food chain. Environmental contamination from pharmaceutical manufacturing and poor sanitation further amplifies resistance.⁴ Bertagnolio and colleagues emphasized the need for global research priorities to address gaps in diagnostics, surveillance, and the discovery of new antimicrobials. They stressed that effective AMR control requires a coordinated “One Health” approach, integrating human, animal, and environmental health.⁵

The clinical consequences of AMR are already evident. Patients with antimicrobial-resistant infections experience higher treatment failure rates, longer hospital stays, and increased mortality. A meta-analysis revealed that multidrug-resistant infections significantly increase mortality risk compared with susceptible infections.⁶ Methicillin-resistant *Staphylococcus aureus*, Carbapenem-resistant *Klebsiella pneumoniae*, Extended-spectrum beta-lactamase (ESBL)-producing organisms are particularly concerning, as they are often resistant to multiple drug classes, with limited available treatment options. The economic burden of AMR is immense. Increased hospitalization, expensive second-line therapies, and the need for infection isolation facilities contribute significantly to healthcare costs. A study reported that if current trends persist, the global cost of AMR could exceed USD 100 trillion by 2050.⁷

Antimicrobial resistance has continued to rise silently across the world despite repeated warnings and stewardship programs for antimicrobial resistance. Many countries have developed and implemented their national action plans on antimicrobial resistance in alignment with the WHO Global Action Plan on AMR. In many low-resource settings, access to reliable diagnostic tools, effective infection control measures, and antimicrobial stewardship programs remains far from adequate.

To tackle this escalating crisis, the global community must act on several urgent priorities. Strengthening laboratory-based surveillance is fundamental; accurate and timely data are essential

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Maria Aslam
Head & Professor, Department of Pathology
Sharif Medical & Dental College, Lahore
E-mail: mariaaslam77@outlook.com

Received: October 15, 2025; Accepted: October 25, 2025

to guide local treatment policies and track emerging resistance patterns. Increased access to rapid diagnostic testing can ensure that antibiotics are used only when truly necessary, preserving their effectiveness for the future. Rigorous infection control measures are vital and should be integrated into every healthcare system's strategy. Governments and health organizations must also provide both financial and regulatory incentives that encourage responsible antibiotic innovation. Without such support, the antibiotic pipeline will continue to decline, leaving clinicians with fewer options to combat increasingly resistant infections.³

Combating it requires a united effort across all sectors: clinicians, pharmacists, policymakers, and the general public. Awareness campaigns should reshape public perception of antibiotics, viewing them not as quick remedies for all illnesses, but as valuable and limited resources that require careful and responsible use. If the world continues on its current trajectory, the consequences will be catastrophic with common infections once again becoming deadly. Strong antibiotic stewardship programs with collective global commitment can still change the course of this crisis. Now it's the time to act decisively together.⁸

REFERENCES

1. World Health Organization. WHO warns of widespread resistance to common antibiotics worldwide. Geneva: WHO. 2025. Available from: <https://www.who.int/news/item/13-10-2025-who-warns-of-widespread-resistance-to-common-antibiotics-worldwide>.
2. GBD 2021 Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance 1990-2021: a systematic analysis with forecasts to 2050. *Lancet*. 2024; 404(10459):1199-226. doi:10.1016/S0140-6736(24)01867-1.
3. American Society of Microbiology. Exploring a one health approach to AMR in Pakistan. *Microcosm Antimicrobial Resistance*. 2023. Available from: <https://asm.org/magazine/2023/fall/exploring-a-one-health-approach-to-amr-in-pakistan>.
4. Centers for Disease Control and Prevention (CDC). Antimicrobial resistance - facts and statistics. Atlanta (GA): CDC. 2025. Available from: <https://www.cdc.gov/antimicrobial-resistance/data-research/facts-stats/index.html>.
5. Bertagnolio S, Dobрева Z, Centner CM, Olaru ID, Dona D, Burzo S, et al. WHO global research priorities for antimicrobial resistance in human health. *Lancet Microbe*. 2024; 5(11):100902. doi:10.1016/S2666-5247(24)00134-4.
6. George NA, Pan D, Silva L, Baggaley RF, Irizar P, Divall P, et al. The prevalence and risk of mortality associated with antimicrobial resistance within nosocomial settings-a global systematic review and meta-analysis of over 20,000 patients. *eClinicalMedicine*. 2025; 87:103384. doi:10.1016/j.eclinm.2025.103384.
7. Ahmed SK, Hussein KQ, Ibrahim RH, Fareeq A, Mahmood KA, Mohamed MG. Antimicrobial resistance: impacts, challenges, and future directions. *Curr Opin Infect Dis*. 2024; 2 (2024):100081. doi:10.1016/j.glmedi.2024.100081.
8. Hashmi A, Ul Haq MI, Malik M, Hussain A, Gajdacs M, Jamshed S. Perceptions of community pharmacists regarding their role in antimicrobial stewardship in Pakistan: A way forward. *Heliyon*. 2023; 9(4):e14843. doi:10.1016/j.heliyon.2023.e14843.



Association of HbA1c Levels with Complexity of Coronary Artery Disease in Diabetic Patients Presenting with Acute Coronary Syndrome

Umer Shafiq, Ismail Ahmed Khan, Sohail Aziz, Kashif ur Rehman, Tayyeb Muhammad, Farman Ullah

ABSTRACT

Objective: To determine the association of Hemoglobin A1c (HbA1c) levels with the complexity of coronary artery disease (CAD) in diabetic patients presenting with acute coronary syndrome (ACS), in terms of higher synergy between percutaneous coronary intervention with taxus and cardiac surgery (SYNTAX) score measured during coronary angiography.

Methodology: This cross-sectional comparative study was conducted at Fauji Foundation Hospital, Rawalpindi from July to September 2025 after approval from the ethical committee of the institution. Patients were divided into two groups. Those with HbA1c $\leq 7\%$ were labeled as group 1 and group 2 had patients with HbA1c $> 7\%$. After informed written consent, 122 confirmed diabetic patients who presented with ACS and underwent coronary angiography were included using non-probability convenience sampling. The SYNTAX score was estimated to determine angiographic disease complexity. The patients with scores of 0-22 were labeled as low risk, 23-32 as intermediate, and > 32 as high-risk CAD. The blood samples of patients were taken and sent for HbA1c levels and fasting lipid profile. Data analysis was carried out with the Statistical Package for the Social Sciences (SPSS) version 26.

Results: When CAD risk categories were compared with HbA1c groups, the majority of the patients in group 1 (34.4%) and group 2 (24.6%) had low risk and intermediate risk CAD, respectively. There was a significant and moderate positive correlation ($r=0.528$) between HbA1c and SYNTAX scores ($p\text{-value}=0.001$). Only the lipid profile showed a significant relationship with CAD risk categories.

Conclusion: The majority of patients with HbA1c $\leq 7\%$ had low-risk CAD, while patients with HbA1c $> 7\%$ had intermediate-risk CAD. The SYNTAX score significantly increased with increasing HbA1c levels. In addition, the patients with intermediate and high-risk CAD also had dyslipidemia.

Keywords: Glycated hemoglobin. Coronary artery disease. Acute coronary syndrome.

INTRODUCTION

Cardiovascular diseases (CVDs) are a major contributor to deaths and disability, leading to 21 million deaths worldwide. Around three quarter of these deaths take place in low- and middle-income countries.¹ In South Asian countries, the burden of CVDs will double in the coming 20 years.² Similarly, diabetes mellitus (DM) is also rising globally, being most common in low- and middle-income countries. The reported prevalence of DM is 537 million across the world. In addition, these diseases are leading to a significant financial burden of healthcare costs. The United Nations Sustainable Development Goal is to decrease the deaths attributed to non-communicable diseases by one-third in these countries.^{1,3}

Acute coronary syndrome (ACS) includes three fatal conditions: ST elevation myocardial infarction (STEMI), non-ST elevation myocardial infarction (NSTEMI), and unstable angina. There are various

predisposing factors of ACS, such as hypertension, obesity, diabetes mellitus, and smoking.⁴ In diabetic patients, coronary artery disease causes a significant proportion of mortality and morbidity.⁵ It has been documented that not only the prevalence of CAD is higher in diabetic patients but DM is also linked with complicated CAD and bad prognosis. This link between DM and CAD is attributed to persistent hyperglycemia in diabetic patients. Hyperglycemia impairs the function of endothelial cells and causes inflammation, and accelerates atherosclerosis.⁶

Hemoglobin A1c (HbA1c) is the standard for monitoring and diagnosis of diabetes mellitus. Some studies have reported the positive relation of HbA1c with the severity of CAD. But others have not found any such association.⁵ Diabetes mellitus is prevalent in Pakistan, reaching a frequency of 17.1%.⁷ Understanding the relation between the two diseases has a great effect on global health owing to the higher prevalence of both diseases. This study was done to determine the association of HbA1c levels with the complexity of CAD indicated by the SYNTAX score in patients presenting with ACS who underwent coronary angiography. This study would provide insight into the possible relation between the two diseases and help in risk stratification for identifying patients at high risk.

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Umer Shafiq
Fellow Interventional Cardiology, Department of Cardiology
Fauji Foundation Hospital, Rawalpindi
E-mail: umer_931849@yahoo.com

Received: September 30, 2025; Accepted: October 28, 2025

METHODOLOGY

This cross-sectional comparative study was conducted at Fauji Foundation Hospital, Rawalpindi after ethical approval (Letter No. 981/RC/FFH/RWP, 30-06-2025) from July to September 2025. Patients were divided into two groups. Those with HbA1c $\leq 7\%$ were labeled as group 1 and group 2 had patients with HbA1c $> 7\%$. The sample size of 61 for each group was calculated using 80% power, 5% margin of error, and the mean SYNTAX scores of 24.33 ± 5.00 & 27.19 ± 6.17 in diabetic patients with HbA1c $\leq 7\%$ & HbA1c $> 7\%$, respectively.⁸ After written informed consent, diabetic patients who presented with ACS and underwent coronary angiography were included using non-probability convenience sampling technique. The exclusion criteria were non-diabetic patients, patients who had previous percutaneous coronary intervention or coronary artery bypass grafting, severe renal or liver diseases, malignancy, and autoimmune disorders.

The medical history & previous laboratory reports of patients were used to confirm diabetes mellitus. The age, body mass index (BMI), and co-morbidities such as hypertension, smoking, dyslipidemia & family history of CAD were noted on a proforma. The blood samples of patients were taken and sent for HbA1c levels and fasting lipid profile. The lipid profile, which includes total cholesterol, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, very low-density lipoprotein (VLDL) cholesterol, and triglycerides, is considered normal when total cholesterol is less than 200 mg/dL, LDL cholesterol is less than 130 mg/dL, HDL cholesterol is greater than 40 mg/dL in men and greater than 50 mg/dL in women, triglycerides are less than 150 mg/dL, and VLDL cholesterol is between 5-30 mg/dL.⁹

The HbA1c levels of $\leq 7\%$ showed good glycemic control and $> 7\%$ was considered suboptimal and often categorized as poor glycemic control.¹⁰ The patients with ACS were diagnosed on suggestive clinical history, raised troponin levels, and electrocardiogram changes.¹¹

The patients then underwent coronary angiography and SYNTAX score was estimated to determine angiographic disease complexity. It was calculated by summing up the score given to individual lesions in the 16 segments of the coronary tree. Patients with $> 50\%$ stenosis in coronary arteries of > 1.5 mm diameter had significant CAD.¹² The range of SYNTAX score was from 0 to > 60 and the higher scores meant more complex CAD. The patients with

scores of 0-22 were labeled as low risk, 23-32 as intermediate and > 32 as high-risk CAD.¹³

STATISTICAL ANALYSIS

Data analysis was carried out using the Statistical Package for the Social Sciences (SPSS) version 26. Numerical and categorical variables were expressed using mean $\pm$ standard deviation and frequency (percentage), respectively. The association between categorical variables like CAD risk categories indicating disease complexity and HbA1c groups was determined using Pearson's Chi-square test. Similar test was used to compare demographic variables and other risk factors with HbA1c groups and CAD risk categories. Pearson's correlation was used to evaluate the correlation between HbA1c levels and SYNTAX scores of the participants. The p-value < 0.05 was considered significant for both tests.

RESULTS

The average age of participants in our study was 54.3 ± 9.35 years. Most of the patients (32.8%) were 51-60 years of age, followed by 41-50 years (28.7%) and > 60 years (27.9%). Most of the patients (77.1%) were females. The mean BMI of the patients was 28.18 ± 4.09 kg/m². Most of the patients (44.3%) were overweight, while 36.8% were obese. Out of 122 patients, 30.3% had hypertension, 41% had a deranged lipid profile, 14.8% were smokers, and 37.7% had a positive family history of CAD. There was no significant difference in demographic variables and co-morbidities between the two groups, showing no significant association of these variables with HbA1c (Table 1).

The statistically significant results of the association of CAD risk categories with HbA1c groups showed that the majority (34.4%) of the patients in group 1 had low risk CAD. However, most (24.6%) of the participants in group 2 had intermediate risk CAD indicating increasing complexity of CAD with increasing HbA1c levels (Table 2).

The mean HbA1c level of the patients was 7.8 ± 1.10 and the mean SYNTAX score was 21.68 ± 6.93 . The Pearson's correlation coefficient was 0.528, indicating a moderate positive correlation between HbA1c & SYNTAX scores (p-value=0.001). This showed that as the HbA1c levels of participants increase, the SYNTAX score, indicating the complexity of CAD, also tends to increase (Figure 1).

When the CAD risk categories were compared with demographic variables and risk factors, only significant association was observed with lipid

profile. The majority (38.5%) of patients with normal lipid profile had low risk CAD indicating lesser complexity of the disease. Most of the patients with intermediate (20.5%) and high risk (9%) CAD

had deranged lipid profiles (Table 3). The other factors, like age groups, gender, BMI, hypertension, smoking, and family history of CAD showed no statistically significant results.

Table 1: Association of Demographic Variables and Risk Factors with HbA1c Groups

Variables		Group 1 (HbA1c ≤7%) (n=61)	Group 2 (HbA1c >7%) (n=61)	Total	p-value
Age Groups (Years)	≤40	8(6.5%)	5(4.1%)	13(10.6%)	0.516
	41-50	20(16.4%)	15(12.3%)	35(28.7%)	
	51-60	18(14.8%)	22(18%)	40(32.8%)	
	>60	15(12.3%)	19(15.6%)	34(27.9%)	
	Total	61(50%)	61(50%)	122(100%)	
Gender	Male	12(9.8%)	16(13.1%)	28(22.9%)	0.389
	Female	49(40.2%)	45(36.9%)	94(77.1%)	
	Total	61(50%)	61(50%)	122(100%)	
BMI (kg/m ²)	Normal (18.5-24.9)	15(12.3%)	8(6.6%)	23(18.9%)	0.077
	Overweight (25-29.9)	29(23.8%)	25(20.5%)	54(44.3%)	
	Obese (≥30)	17(13.9%)	28(22.9%)	45(36.8%)	
	Total	61(50%)	61(50%)	122(100%)	
Clinical Presentation	NSTEMI	34(27.9%)	40(32.8%)	74(60.7%)	0.266
	STEMI	27(22.1%)	21(17.2%)	48(39.3%)	
	Total	61(50%)	61(50%)	122(100%)	
Hypertension	Hypertensive	17(13.9%)	20(16.4%)	37(30.3%)	0.554
	Non-Hypertensive	44(36.1%)	41(33.6%)	85(69.7%)	
	Total	61(50%)	61(50%)	122(100%)	
Lipid Profile	Deranged	24(19.7%)	26(21.3%)	50(41%)	0.712
	Normal	37(30.3%)	35(28.7%)	72(59%)	
	Total	61(50%)	61(50%)	122(100%)	
Smoking	Smoker	10(8.2%)	8(6.6%)	18(14.8%)	0.609
	Non-Smoker	51(41.8%)	53(43.4%)	104(85.2%)	
	Total	61(50%)	61(50%)	122(100%)	
Family History of CAD	Positive	21(17.2%)	25(20.5%)	46(37.7%)	0.454
	Negative	40(32.8%)	36(29.5%)	76(62.3%)	
	Total	61(50%)	61(50%)	122(100%)	

Table 2: Association of CAD Risk Categories with HbA1c Groups

CAD Risk Categories (SYNTAX Score)	HbA1c Groups		Total	p-value
	Group 1 (HbA1c ≤7%)	Group 2 (HbA1c >7%)		
Low Risk (0-22)	42(34.4%)	19(15.6%)	61(50%)	0.001*
Intermediate Risk (23-32)	14(11.5%)	30(24.6%)	44(36.1%)	
High Risk (>32)	5(4.1%)	12(9.8%)	17(13.9%)	
Total	61(50%)	61(50%)	122(100%)	

*Significant p-value

Table 3: Association of CAD Risk Categories with Lipid Profile

Lipid Profile	CAD Risk Categories (SYNTAX Score)			Total	p-value
	Low Risk (0-22) n=61	Intermediate Risk (23-32) n=44	High Risk (>32) n=17		
Deranged	14(11.5%)	25(20.5%)	11(9%)	50(41%)	0.0002*
Normal	47(38.5%)	19(15.6%)	6(4.9%)	72(59%)	
Total	61(50%)	44(36.1%)	17(13.9%)	122(100%)	

*Significant p-value




Figure 1: Scatter Plot showing a Positive Association between HbA1c and SYNTAX Score

DISCUSSION

The variation in blood glucose levels strongly influences the development of CAD. Hemoglobin A1c is a prognostic indicator of glycemic control in diabetic patients and can predict the complexity of CAD. Therefore, it is mandatory to monitor HbA1c levels continuously for risk assessment of patients.^{14,15}

The average age of patients in our study was 54.3 ± 9.35 years, with the majority (77.1%) of females. However, the age groups and gender did not differ significantly when compared with HbA1c groups and CAD risk categories. In a study done by Rashid et al., the mean age was 58.77 ± 18.24 years with 60% males. In contrast to our results, the mean age of their participants had a statistically significant association ($p=0.06$) with CAD severity where increased severity was shown by older ages. The gender didn't show any such association.¹⁶ Qadir et al. observed that 56.6% of their patients were males and the average age was 57.54 ± 3.47 years.¹⁷ A study from China showed that the average age of their participants was 65 years with 64.3% females. The age and gender didn't differ significantly when compared with adverse cardiac events indicating the complexity and severity of CAD in diabetic patients.¹⁸ The average age was 57.6 ± 9.5 years in another study with 65% males.¹⁹ Patients had an average age of 54 ± 10.2 years with 79.8% males in another study conducted in Karachi, Pakistan. But neither age nor gender had any statistically

significant association with the prediction of CAD severity.²⁰

Our results revealed that the mean BMI of the patients was 28.18 ± 4.09 kg/m². Most of the patients (44.3%) were overweight, while 36.8% were obese. There was no significant association between BMI and HbA1c groups and CAD risk categories. Jiao et al. showed that the mean BMI of participants was 24.9 ± 2.4 kg/m² and BMI and disease severity/complexity in diabetic patients had no significant association ($p>0.05$).¹⁷ Another study reported that BMI had no statistically significant role in the prediction of CAD severity.²⁰ The risk factors profile of the current study showed that 30.3% of the patients were hypertensive, 14.8% were smokers, 41% had a deranged lipid profile, and 37.7% had a family history of CAD. None of these risk factors showed a statistically significant association when compared with HbA1c groups or CAD risk categories except lipid profile. The majority (38.5%) of patients with normal lipid profile had low risk CAD indicating lesser complexity, while most of the patients with intermediate (20.5%) and high risk (9%) CAD had deranged lipid profiles. Rashid et al. observed in their study that there were 73.77% hypertensive, 44.89% smokers, and 56.88% patients with dyslipidemia. These risk factors showed no significant relationship with the severity of CAD.¹⁶ Similar to our results, Garg et al. observed that a deranged lipid profile was significantly related to CAD severity.¹⁹ The most common risk factor in a

study was smoking (44.5%), followed by hypertension (42%), family history of CAD (37%), and dyslipidemia (10%). These risk factors didn't serve as significant predictors of CAD severity.²⁰

Our results revealed that most of the ACS patients (60.7%) presented with NSTEMI. More than half of the patients presented with NSTEMI (52.1%), followed by STEMI (25.2%) and unstable angina (22.7%) in another study done by Habib et al. Similar to our results, clinical presentation wasn't related to CAD complexity.²⁰ However, a previous study observed that patients with STEMI presentation had more complex disease indicated by higher SYNTAX scores ($p<0.05$).²¹

The current study reported statistically significant results of the association of CAD risk categories with HbA1c groups. The majority (34.4%) of the patients in group 1 had low risk CAD. However, most (24.6%) of the participants in group 2 had intermediate risk CAD indicating increasing complexity of CAD with higher HbA1c levels. Another study showed that the patients with higher HbA1c levels had significantly increased complexity/severity of CAD indicated by higher Gensini scores.¹⁶ Similar results were shown by another study, which revealed that HbA1c was a significant predictor of the severity of coronary artery disease and major adverse cardiac events.¹⁸ Garg et al. revealed a significant association between HbA1c in diabetic patients and severity of CAD, where 41.7% of the patients with HbA1c $>10.5\%$ had triple vessel CAD.¹⁹ Our study showed a significant and moderate positive correlation of HbA1c and SYNTAX scores. Similar results were reported in a study by Xu et al., where HbA1c and SYNTAX scores were significantly & positively correlated in diabetic patients.⁸ The study by Dar et al. also found a significant correlation between HbA1c and Gensini scores showing increasing complexity of CAD with higher HbA1c levels.²² Habib et al. conducted a study on non-diabetic patients with ACS and did not find any significant correlation between HbA1c and SYNTAX scores.²⁰

CONCLUSION

The majority of patients with HbA1c $\leq 7\%$ had low-risk CAD, while patients with HbA1c $>7\%$ had intermediate-risk CAD. Our study also found a significant, moderate positive correlation between HbA1c and the complexity of coronary artery disease (indicated by SYNTAX score) in diabetic patients presenting with ACS. The SYNTAX score significantly increased with increasing HbA1c levels. In addition, the patients with intermediate and

high-risk CAD indicated by higher SYNTAX scores also had dyslipidemia.

LIMITATIONS & RECOMMENDATIONS

This study had a few limitations: cross-sectional design, non-probability convenience sampling technique, and single-institution research. The study did not evaluate the association of higher HbA1c in diabetic patients with complications of CAD, such as cardiac mortality and major adverse cardiovascular events. In the future, multi-centered studies should be conducted with follow-up of the patients to assess long-term complications.

Diabetic patients with higher HbA1c levels should be considered at higher cardiovascular risk as higher HbA1c is linked with higher SYNTAX scores, indicating complex CAD. The need for strict glycemic control in diabetic patients should be emphasized to reduce cardiovascular risk.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

U.S: Assisted in patient recruitment and data acquisition, analyzed data, and drafted the manuscript.

I.A.K: Contributed to study design, data interpretation, and critical revision.

S.A: Conceived and designed the study.

K.R: Performed statistical analysis and literature review.

T.M: Assisted in data collection and manuscript formatting.

F.U: Supervised the study, validated results, and approved the final manuscript.

REFERENCES

1. Di Cesare M, Perel P, Taylor S, Kabudula C, Bixby H, Gaziano TA, et al. The heart of the world. *Glob Heart*. 2024; 19(1):11. doi:10.5334/gh.1288.
2. Panhwar S, Memon SH, Soomro MK, Soomro TH, Memon AG. Association of HbA1c level with extent of acute myocardial infarction and short term mortality. *JIMDC*. 13(2):2024. doi:10.35787/jimdc.v13i2.1223.
3. Hossain MJ, Al-Mamun M, Islam MR. Diabetes mellitus, the fastest growing global public health concern: early detection should be focused. *Health Sci Rep*. 2024; 7(3):e2004. doi:10.1002/hsr2.2004.
4. Khan FR, Ali J, Ullah R, Hassan Z, Khattak S, Lakhta G, et al. Relationship between high glycated hemoglobin and severity of coronary artery disease in type II diabetic patients hospitalized with acute coronary syndrome. *Cureus*. 2021; 13(3):e13734. doi:10.7759/cureus.13734.
5. Mohamed AA, Abd El-Rahman YH, Sabry SM, Abdel Kader AS. Hemoglobin A1c in non-diabetic NSTEMI patients as a predictor of severity of coronary artery disease. *Benha Med J*. 2024; XX(XX). doi:10.21608/bmfj.2024.327152.2225.

6. Albashir AAD, Elawad OAMA, Khougali Mohamed H. The use of glycosylated hemoglobin (HbA1c) as a predictor of the severity of acute coronary syndrome among diabetic patients. *Ir J Med Sci.* 2021; 190(2):609-14. doi:10.1007/s11845-020-02341-0.
7. Wasay A, Shaikh A, Kumari M, Ishaq M, Ali M, Sadik A, et al. Frequency of undiagnosed diabetes in patient presenting with acute coronary syndrome at tertiary care hospital of Karachi. *IJHR.* 2025; 3(1):631-9. doi:10.71000/q03q8b42.
8. Xu S, Zhou B, Zhou B, Zheng J, Xu Q, Wang B, et al. The level of HbA1c evaluates the extent of coronary atherosclerosis lesions and the prognosis in diabetes with acute coronary syndrome. *Comput Math Methods Med.* 2022; 2022:7796809. doi:10.1155/2022/7796809.
9. Pancholia AK, Kabra NK, Gupta R. Laboratory evaluation of lipid parameters in clinical practice. *Indian Heart J.* 2024; 76(Suppl 1):S29-32. doi: 10.1016/j.ihj.2024.02.002.
10. Adjei SK, Adjei P, Nkrumah PA. Poor glycemic control and its predictors among type 2 diabetes patients: insights from a single-centre retrospective study in Ghana. *Health Sci Rep.* 2025; 8(3):e70558. doi:10.1002/hsr2.70558.
11. Bergmark BA, Mathenge N, Merlini PA, Lawrence-Wright MB, Giugliano RP. Acute coronary syndromes. *Lancet.* 2022; 399(10332):1347-58. doi:10.1016/S0140-6736(21)02391-6.
12. Derimay F, Finet G, Rioufol G. Coronary artery stenosis prediction does not mean coronary artery stenosis obstruction. *Eur Heart J.* 2021; 42(42):4401. doi:10.1093/eurheartj/ehab332.
13. Khawaja AZ, Khalil M, Shahzad SK, Yousaf A, Anwar M, Ahmed I, et al. Association of severity of coronary artery disease with SYNTAX score and ankle brachial index. *PAFMJ.* 2024; 74. doi:10.51253/pafmj.v74i-SUPP-1-10579
14. Singh B, Agarwal D, Dahiya A, Saxena GN. Evaluation of the association between glycated hemoglobin levels and severity of coronary artery disease in non-diabetic patients. *Heart India.* 2021; 9(3):184. doi:10.4103/heartindia.heartindia_99_21.
15. Chen J, Yin D, Dou K. Intensified glycemic control by HbA1c for patients with coronary heart disease and type 2 diabetes: a review of findings and conclusions. *Cardiovasc Diabetol.* 2023; 22(1):146. doi:10.1186/s12933-023-01875-8.
16. Rashid A, Saeed MS, Ghouse MA, Kunju AK, Umar N, Asif AE, et al. Association between glycated hemoglobin and severity of coronary artery disease in type 2 diabetic patients with myocardial infarction. *Cureus.* 2025; 17(3):e81521. doi:10.7759/cureus.81521.
17. Qadir M, Khan MSN, Khan HR, Ullah N, Ahmed M, Rashid HMF. Association between HbA1c and coronary artery disease severity in patients of diabetes mellitus type 2 presented as acute coronary artery syndrome and non-diabetic. *PJMHS.* 2022; 16(09):285-7. doi:10.53350/pjmhs.22169285.
18. Jiao X, Zhang Q, Peng P, Shen Y. HbA1c is a predictive factor of severe coronary stenosis and major adverse cardiovascular events in patients with both type 2 diabetes and coronary heart disease. *Diabetol Metab Syndr.* 2023; 15(1):50. doi:10.1186/s13098-023-01015-y.
19. Garg Y, Marthandam Manickandan AP, Ghori HK, Rameshabu S, Mohammed I, Mohamed S, et al. Analysis of angiographic patterns in acute coronary syndrome patients with diabetes mellitus: correlation with HbA1c levels. *Cureus.* 2024; 16(10):e72028. doi:10.7759/cureus.72028.
20. Habib S, Ullah SZ, Saghir T, Syed Muhammad A, Ud Deen Z, Naseeb K, et al. The association between hemoglobin A1c and the severity of coronary artery disease in non-diabetic patients with acute coronary syndrome. *Cureus.* 2020; 12(1):e6631. doi:10.7759/cureus.6631.
21. Abbaszadeh S, Rafati S, Mamikhani D, Emami M, Shahabi N. Predictive power of glycated hemoglobin in detecting severity of coronary artery disease in non-diabetic patients: a cross-sectional study in Southern Iran. *ARYA Atheroscler.* 2024; 20(5):15-24. doi:10.48305/arya.2024.42463.2936.
22. Dar MI, Beig JR, Jan I, Shah TR, Ali M, Rather HA, et al. Prevalence of type 2 diabetes mellitus and association of HbA1c with severity of coronary artery disease in patients presenting as non-diabetic acute coronary syndrome. *Egypt Heart J.* 2020; 72: 66. doi:10.1186/s43044-020-00101-0.



Efficacy of Lacosamide in Patients with Diabetic Neuropathy

Awais Akram, Shahzad Alam Khan, Umair Ahmad

ABSTRACT

Objective: To compare the efficacy of lacosamide versus placebo in patients with diabetic neuropathy in terms of reduction of the mean pain score.

Methodology: This randomized controlled trial (RCT) was conducted at the Department of Medicine, Nishtar Hospital, Multan from April to August 2025. After informed written consent, 60 patients with diabetic neuropathy were included and their baseline pain levels at recruitment were assessed using the visual analogue scale (VAS). The patients were enrolled using non-probability convenience sampling technique and subsequently randomized equally through a lottery method. Randomization was used to assign patients to either group A (lacosamide) or group B (placebo). Participants were administered either lacosamide (LCM; brand name atcomid) or a placebo that contained microcrystalline cellulose, which was commercially available and widely used in clinical trials involving diabetics. Both placebo & lacosamide were started at 100 mg daily and escalated to 400 mg/day over four weeks. Both were maintained at 400 mg for 12 weeks. The VAS was used to measure baseline and post-treatment pain scores. The variables of age, gender, duration of diabetes, obesity (body mass index ≥ 30 kg/m²), and diabetes control were documented. Statistical Package for the Social Sciences (SPSS) version 26 was used for the analysis of the data.

Results: The age distribution of the study population was statistically similar in the two groups (46.67 $\pm$ 9.66 years vs. 49.30 $\pm$ 8.38 years; $p=0.264$). Majority of the participants were males in both groups (66.7% vs. 56.7%). The baseline VAS values were almost similar across the groups (group A: 7.40 $\pm$ 1.96 vs. group B: 6.87 $\pm$ 2.01; $p=0.31$). The lacosamide group showed a notable decrease in post-treatment pain scores as compared to placebo (3.90 $\pm$ 1.92 vs. 6.73 $\pm$ 1.72; $p<0.001$).

Conclusion: Lacosamide provided a statistically and clinically significant reduction in pain intensity among patients with diabetic neuropathy compared to placebo. The marked decrease in post-treatment VAS scores highlighted its potential as an effective adjunctive option for managing neuropathic pain in this population.

Keywords: Diabetic neuropathy. Diabetes mellitus. Small fiber neuropathy. Visual analog scale. Lacosamide.

INTRODUCTION

A significant public health challenge of this century is Diabetes mellitus. Its extended duration leads to a multitude of macro and microvascular complications that impact nearly all organ systems.¹ One in five individuals diagnosed with diabetes experiences chronic diabetic painful neuropathy (DPN). Distal symmetric polyneuropathy, which represents 80-90% of diabetic neuropathies, involves both small and large fiber damage. It is typically characterized by sensory disturbances that begin in the feet, progress upward to the calves, and, in advanced stages, extend to the upper limbs.² Damage to large nerve fibers leads to paresthesia, sensory loss, and muscle weakness, whereas small fiber damage is linked to pain, anesthesia, foot ulcers, and autonomic dysfunction.³ Painful diabetic peripheral neuropathy significantly

reduces quality of life in individuals with diabetes and, if left unrecognized and untreated, increases morbidity and mortality through non-traumatic amputations. Neuropathic symptoms, however, may be minimized with proper blood glucose control, lifestyle modifications, dietary management, and regular follow-up.⁴

Pharmacological approaches to managing DPN encompass a range of options, including antiepileptics like pregabalin & gabapentin, opioids, selective serotonin reuptake inhibitors, norepinephrine reuptake inhibitors, and tricyclic antidepressants (TCA). However, it is essential to note that these treatments may not yield optimal results for all individuals.⁵ Furthermore, certain medications may exhibit significant negative consequences when used over prolonged periods.³

Lacosamide (LCM) is a newly recognized antiepileptic drug that demonstrated analgesic and neuroprotective properties across various studies. It works by blocking voltage-gated sodium channels in a slow inactivating manner and is linked to side effects like nausea and dizziness. Research conducted within Western populations has shown that LCM effectively alleviates neuropathic pain and is generally well tolerated in cases of DPN.⁶

Many foreign studies have examined lacosamide in diabetic neuropathy. Due to variations in glycemic

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Awais Akram
Postgraduate Resident, Department of Medicine
Nishtar Hospital, Multan
E-mail: awais.szmc@gmail.com

Received: September 25, 2025; Accepted: October 27, 2025

management patterns, comorbidity burden, and medication metabolism, the effectiveness and tolerance of lacosamide may differ among local and international populations. There is a paucity of local evidence on the effect of lacosamide in painful diabetic neuropathy, where diabetes is among the most common diseases, and neuropathic consequences cause significant morbidity. Thus, this study fills a significant gap by comparing lacosamide to a placebo in Pakistani patients to enhance local guidelines and improve patient outcomes.

METHODOLOGY

This randomized controlled trial was conducted at the Department of Medicine, Nishtar Hospital, Multan from April to August 2025. The study was approved by the institutional ethical board (Letter No. 3607/NMU, 10-03-2025) and registered with the Iranian Trial Registry (IRCT20250211064704N1; 21-05-2025). The sample size was calculated via OpenEpi software with 80% power, 95% confidence level, and based on mean post-treatment pain scores of 3.39 ± 1.94 in the lacosamide group and 6.35 ± 1.52 in the placebo group.⁷ After informed written consent, 60 patients were included in the study and they were divided into two groups (30 in each). Male and female patients, aged between 20 to 60 years who had been diagnosed with diabetes mellitus for a minimum of five years and developed diabetic neuropathy were included. Patients were excluded if they had cardiovascular disease, renal impairment, elevated liver enzymes that were twice the normal range. Additionally, those receiving treatment with any painkiller like non-steroidal anti-inflammatory drugs, tramadol, TCA, mexiletine hydrochloride, lidocaine patch, or opioids were also excluded. Pregnant and breastfeeding mothers were also not included. The patients were enrolled using non-probability convenience sampling technique and subsequently randomized equally through a lottery method. A lottery employing sealed, opaque envelopes assigned patients to either group A (lacosamide) or group B (placebo).

Diabetic neuropathy was assessed clinically with history and physical examination for small and large fiber neuropathy. Age, gender, duration of diabetes, obesity (body mass index ≥ 30 kg/m²), and diabetes control were documented. Diabetes control was evaluated by HbA1c results. HbA1c levels of all the patients were tested from the same laboratory. Patients exhibiting HbA1c values of 7% or lower had controlled diabetes, while those with values of more than 7% were classified as having uncontrolled

diabetes. Participants were administered either lacosamide (LCM; brand name atcomid) or a placebo that contained microcrystalline cellulose, which was commercially available and widely used in clinical trials involving diabetics. Both placebo & lacosamide were given as oral tablets, starting from 100 mg/day during the first week and escalated to a maximum of 400 mg per day at the end of four weeks, with a weekly increase of 100 mg each week. After achieving the maximum dose of 400 mg/day as 200 mg BD (twice a day), the patients were maintained on this dosage for twelve weeks. Pain levels were assessed using the VAS at baseline and after a three-month treatment period with 400 mg/day dose. Pain scores ranged from 0 to 10 (0 as no pain, and 10 as extreme pain). In accordance with the established protocol, an evaluator, unaware of the treatment allocation, assessed the patients' pain levels at baseline and after the therapeutic intervention.

STATISTICAL ANALYSIS

The data analysis was performed using Statistical Package for the Social Sciences (SPSS) version 26. The mean $\pm$ standard deviations were presented for age, duration of diabetes, HbA1c level, and pain scores. The frequencies & percentages of gender, obesity, and both controlled & uncontrolled diabetes were documented. Categorical data were analyzed using the Chi-square test and an independent t-test was used to compare means of quantitative variables between two groups. Paired t-test was applied to compare mean pain scores at baseline and post-treatment (3 months) within the groups. A p-value of <0.05 was considered significant.

RESULTS

The mean age of patients was 46.67 ± 9.66 years in group A and 49.30 ± 8.38 years in group B, and most of the patients in both groups were more than 40 years old. The male demographic was more prevalent in both groups. The mean duration of diabetes mellitus in groups A and B was 22.90 ± 13.08 years and 23.50 ± 13.94 years, respectively. Less than half in both groups were obese (40% vs. 43.3%). However, both of the groups showed no statistically significant difference in terms of distributions of age, gender, obesity, and duration of diabetes mellitus. The two groups also showed no significant difference in terms of mean HbA1c levels ($5.97 \pm 2.01\%$ vs. $6.71 \pm 1.77\%$) (Table 1).

The baseline VAS values for pain were similar across the groups, with group A scoring 7.40 ± 1.96

and group B scoring 6.87 ± 2.01 ($p=0.31$). Group A experienced a significant decrease in pain score to 3.90 ± 1.92 , whereas group B exhibited no significant difference after treatment ($p=0.736$). A statistically significant difference in post-treatment scores was found between the study groups ($p < 0.001$) (Table 2).

Regarding age, gender, obesity, and diabetes control, mean VAS score was significantly lower in Lacosamide group ($p < 0.001$). The duration of diabetes did not significantly influence treatment response in terms of VAS improvement ($p > 0.05$) (Table 3).

Table 1: Demographic and Baseline Characteristics of the Study Groups

Characteristics		Group A (Lacosamide) Frequency & Percentage	Group B (Placebo) Frequency & Percentage	p-value
Age Groups (Years)	20-40	9(30%)	5(16.7%)	0.22
	41-60	21(70%)	25(83.3%)	
Gender	Male	20(66.7%)	17(56.7%)	0.43
	Female	10(33.3%)	13(43.3%)	
Duration of Diabetes Mellitus (Years)	≤ 10	8(26.7%)	9(30%)	0.77
	> 10	22(73.3%)	21(70%)	
Obesity	Yes	12(40%)	13(43.3%)	0.79
	No	18(60%)	17(56.7%)	
Diabetes Control	Controlled	20(66.7%)	14(46.7%)	0.12
	Un-Controlled	10(33.3%)	16(53.3%)	

Table 2: Baseline and Post-Treatment Pain Score of the Study Groups

VAS score	Group A (Lacosamide)	Group B (Placebo)	p-value
Baseline Pain Score (Mean $\pm$ SD)	7.40 ± 1.96	6.87 ± 2.01	0.31
Post-Treatment Pain Score (Mean $\pm$ SD)	3.90 ± 1.92	6.73 ± 1.72	$< 0.001^*$
p-value	0.001*	0.736	

*Significant p-value

Table 3: Post-Stratification Analysis of Post-Treatment VAS Scores in Groups A & B

Variables		Group A (Lacosamide) Mean $\pm$ SD	Group B (Placebo) Mean $\pm$ SD	p-value
Age Groups (Years)	20-40	4.44 ± 2.06	7.00 ± 2.12	$< 0.001^*$
	41-60	3.67 ± 1.85	6.68 ± 1.68	$< 0.001^*$
Gender	Male	3.90 ± 1.84	6.47 ± 1.88	$< 0.001^*$
	Female	3.90 ± 2.18	7.08 ± 1.49	$< 0.001^*$
Duration of Diabetes Mellitus (Years)	≤ 10	4.00 ± 1.69	3.86 ± 2.03	0.77
	> 10	6.44 ± 1.81	6.86 ± 1.71	0.36
Obesity	Yes	3.25 ± 1.54	7.23 ± 1.42	$< 0.001^*$
	No	4.33 ± 2.05	6.35 ± 1.86	0.0002*
Diabetes Control	Controlled	3.95 ± 1.72	7.29 ± 1.68	$< 0.001^*$
	Un-controlled	3.80 ± 2.25	6.25 ± 1.65	$< 0.001^*$

*Significant p-value

DISCUSSION

In Pakistan, diabetes stands as the leading cause of peripheral neuropathy. The occurrence of peripheral neuropathy in individuals with diabetes in Pakistan exhibits considerable variation, with rates spanning from 16.30% to 79.50%.⁸

In our study, the mean age of the patients was 46.67 ± 9.66 years in group A and 49.30 ± 8.38 years in group B with a majority of males. A study reported that most of the patients with diabetic neuropathy were males (61.8%) and above 60 years old (85.6%).⁹ Hussain et al. observed that age above 50 years was significantly associated with DNP.⁸ A study conducted at Sheikh Zayed Hospital, Lahore, reported a 20% prevalence of DNP in patients aged ≥ 60 years, compared to 8.9% in those aged ≤ 35 years.¹⁰ The majority of patients presenting with diabetic neuropathy had more than 10 years of duration since diagnosis, and the mean duration of diabetes mellitus in groups A and B was 22.90 ± 13.08 and 23.50 ± 13.94 years, respectively. Consistent with our findings, a nationwide study depicted that over 10 years after the diagnosis of diabetes was associated with painful DNP. Moreover, the presence of other co-morbid conditions like hypertension and ischemic heart disease was also related to the early development of diabetic neuropathy.¹¹ Bondar et al. also stated that DNP was the most prevalent microvascular complication in individuals with diabetes, affecting more than half of patients after 20 years of disease progression.¹²

Despite being on hypoglycemic medications, 43.3% of patients in the current study had uncontrolled diabetes, indicated by HbA1C levels above 7%. Previous literature emphasized that chronic hyperglycemia increased the risk of peripheral diabetic neuropathy.¹³ Nozawa et al. also reported that higher 3-year mean HbA1c levels showed a significant association with the occurrence of DPN (adjusted odds ratio: 1.23; 95% CI: 1.06-1.42).⁹ Another study observed that increasing HbA1c levels were significantly associated with the occurrence of DNP ($p < 0.05$).¹¹ Hepsen et al. found that even HbA1c levels of prediabetics showed a significant positive correlation ($r=0.188$, $p=0.014$) with VAS score.¹⁴

Despite the effectiveness of traditional treatments like pregabalin, duloxetine, or tricyclic antidepressants, neuropathic pain continued to be a considerable contributor to morbidity and disability.^{13,15} When the mean post-treatment pain scores were evaluated, the lacosamide group exhibited a significant reduction in pain scores from

7.40 ± 1.96 to 3.90 ± 1.92 ($p < 0.001$), compared to the placebo group in our study. A study from Sindh, Pakistan reported that patients in the lacosamide group had a mean pre-trial pain score of 6.59 ± 1.95 , which declined to 3.39 ± 1.94 . In comparison, the placebo group showed a baseline score of 6.71 ± 1.89 , decreasing only to 6.35 ± 1.52 after the 20-week maintenance phase ($p < 0.0001$).⁷

In a recent multicenter trial to study the effect of lacosamide on peripheral neuropathic pain, lacosamide reduced the pain intensity by up to 50% in patients with peripheral neuropathic pain.¹⁶ In a meta-analysis, LCM was reported to be more effective than placebo in reducing the pain of DPN, but showed inferior efficacy when compared to 1st line drugs like duloxetine and pregabalin. However, it was more tolerable and had fewer side effects.¹⁷ In a study conducted in Swat, Riaz et al. combined LCM with pregabalin and reported that the pregabalin group demonstrated a mean post-treatment score of 3.07 ± 1.32 , whereas the combination of pregabalin and lacosamide achieved a lower mean score of 2.30 ± 1.16 ($p < 0.05$). Treatment effectiveness was observed in 58.2% of the pregabalin group compared to 80.6% in the combination group ($p < 0.05$).¹⁸ In a recent study on the Indian population, Lacosamide significantly reduced pain (numeric rating score: 7.7 to 2.50) compared to pregabalin (7.6 to 4.27) ($p < 0.001$), indicating greater effectiveness.¹⁹

CONCLUSION

Lacosamide showed significantly higher efficacy in reducing pain intensity indicated by lower VAS scores among patients with diabetic neuropathy compared to placebo. When compared with placebo, lacosamide significantly lowered pain scores across age, gender, diabetes control, and obesity groups. The marked decrease in post-treatment VAS scores highlighted its potential as an effective adjunctive option for managing neuropathic pain in diabetic patients.

LIMITATIONS & RECOMMENDATIONS

Our study had a few limitations, including a single-centered study and a small sample size. Moreover, a fixed dose of lacosamide was studied. Further studies are recommended on lacosamide at various doses, with a special emphasis on its side effects and tolerability. Studies on combination treatments are also recommended for the future.

Conflict of interest: None.

Source of funding: None.

Authors Contributions:

A.A: Conception of the idea and data collection

S.A.K: Data collection and analysis

U.A: Manuscript writing and proofreading

REFERENCES

1. Tomic D, Shaw JE, Magliano DJ. The burden and risks of emerging complications of diabetes mellitus. *Nat Rev Endocrinol.* 2022; 18(9):525-39. doi:10.1038/s41574-022-00690-7.
2. Jensen TS, Karlsson P, Gylfadottir SS, Andersen ST, Bennett DL, Tankisi H, et al. Painful and non-painful diabetic neuropathy, diagnostic challenges and implications for future management. *Brain.* 2021; 144(6):1632-45. doi:10.1093/brain/awab079.
3. Marshall A, Alam U, Themistocleous A, Calcutt N, Marshall A. Novel and emerging electrophysiological biomarkers of diabetic neuropathy and painful diabetic neuropathy. *Clin Ther.* 2021; 43(9):1441-56. doi:10.1016/j.clinthera.2021.03.020.
4. Say B, Ergun U, Yildiz A, Alpua M, Arikan Durmaz S, Turgal E. Detecting pain severity with full cup test in painful diabetic peripheral neuropathy. *Agri.* 2020; 32(2):85-90. English. doi:10.14744/agri.2019.82653.
5. Pop-Busui R, Ang L, Boulton AJM, Feldman EL, Marcus RL, Mizokami-Stout K, et al. Diagnosis and treatment of painful diabetic peripheral neuropathy. Arlington (VA): American Diabetes Association. 2022; 2022(1):1-32. doi: 10.2337/db2022-01.
6. Munoz-Vendrell A, Teixidor S, Sala-Padro J, Campoy S, Huerta-Villanueva M. Intravenous lacosamide and phenytoin for the treatment of acute exacerbations of trigeminal neuralgia: a retrospective analysis of 144 cases. *Cephalalgia.* 2022; 42(10):1031-8. doi:10.1177/03331024221092435.
7. Rehman MU, Ahmed Z, Mujtaba G, Ali A, Shah S, Singh J. Role of lacosamide in diabetic neuropathy: an experience at a tertiary care hospital in rural Sindh. *PJMHS.* 2022; 16(11):293-5. doi:10.53350/pjmhs 20221611293.
8. Hussain A, Ahmad I, Khan MN, Zar A, Shahzad S. Peripheral neuropathy among patients with type 2 diabetes using the Michigan neuropathy screening instrument. *J Postgrad Med Inst.* 2021; 35(1):7-11. doi:10.54079/jpmi.35.1.2735.
9. Nozawa K, Ikeda M, Kikuchi S. Association between HbA1c levels and diabetic peripheral neuropathy: a case-control study of patients with type 2 diabetes using claims data. *Drugs Real World Outcomes.* 2022; 9(3):403-14. doi:10.1007/s40801-022-00309-3.
10. Perveen W, Ahsan H, Rameen Shahzad, Fayyaz S, Zaif A, Paracha MA, et al. Prevalence of peripheral neuropathy, amputation, and quality of life in patients with diabetes mellitus. *Sci Rep.* 2024; 14(1):14430. doi:10.1038/s41598-024-65495-2.
11. Li C, Wang W, Ji Q, Ran X, Kuang H, Yu X, et al. Prevalence of painful diabetic peripheral neuropathy in type 2 diabetes mellitus and diabetic peripheral neuropathy: a nationwide cross-sectional study in mainland China. *Diabetes Res Clin Pract.* 2023; 198:110602. doi:10.1016/j.diabres.2023.110602.
12. Bondar A, Popa AR, Papanas N, Popoviciu M, Vesa CM, Sabau M, et al. Diabetic neuropathy: a narrative review of risk factors, classification, screening and current pathogenic treatment options (Review). *Exp Ther Med.* 2021; 22(1):690. doi:10.3892/etm.2021.10122.
13. Tesfaye S, Sloan G, Petrie J, White D, Bradburn M, Julious S, et al. Comparison of amitriptyline supplemented with pregabalin, pregabalin supplemented with amitriptyline, and duloxetine supplemented with pregabalin for the treatment of diabetic peripheral neuropathic pain (OPTION-DM): a multicentre, double-blind, randomised crossover trial. *Lancet.* 2022; 400(10353):680-90. doi:10.1016/S0140-6736(22)01472-6.
14. Hepsen S, Keskin DB, Cakal E. The assessment of neuropathic pain in patients with prediabetes. *Prim Care Diabetes.* 2023; 17(1):33-7. doi:10.1016/j.pcd.2022.11.004.
15. Roikjer J, Borbjerg MK, Andresen T, Giordano R, Hviid CVB, Morch CD, et al. Diabetic peripheral neuropathy: emerging treatments of neuropathic pain and novel diagnostic methods. *J Diabetes Sci Technol.* 2024:19322 968241279553. doi:10.1177/19322968241279553.
16. Carmland ME, Kreutzfeldt MD, Holbech JV, Brask-Thomsen PK, Kroigard T, Hansen PN, et al. The effect of lacosamide in peripheral neuropathic pain: a randomized, double-blind, placebo-controlled, phenotype-stratified trial. *Eur J Pain.* 2024; 28(1):105-19. doi:10.1002/ejp.2165.
17. Jingxuan L, Litian M, Jianfang F. Different drugs for the treatment of painful diabetic peripheral neuropathy: a meta-analysis. *Front Neurol.* 2021; 12:682244. doi:10.3389/fneur. 2021.682244.
18. Riaz M, Khalid, Ali S, Ali A, Shah MA, Rehanullah, et al. Efficacy of pregabalin alone and in combination with lacosamide in pain management of diabetic neuropathy. *J Saidu Med Coll.* 2025; 15(3):390-7. Available from: <https://www.jsmc.pk/index.php/jsmc/article/view/1129/698>.
19. Kalpana R, Robert WS, Juzerehaji F, Venkatachalapathy R, Jagadeesan M. A comparison of the effectiveness of lacosamide and pregabalin in treating patients diagnosed with neuropathic pain: a randomized, controlled, open-label pilot trial. *Naunyn Schmiedeberg Arch Pharmacol.* 2025; 12:682244. doi:10.1007/s00210-025-04474-4.



Intestinal Parasitic Infections among Children and Young Adolescents Visiting Provincial Headquarter Hospital in Gilgit, Pakistan and their Associated Factors

Uzma Ali, Sadaf Nasir, Adnan Qureshi, Aqsa Aslam, Namra Yunus, Zakia Wahid

ABSTRACT

Objective: To detect the frequency and species distribution of intestinal parasites among children and young adolescents visiting a provincial Headquarter Hospital in Gilgit, Pakistan and determine their association with socio-demographic and environmental factors

Methodology: This descriptive, cross-sectional study was done among patients visiting the Provincial Headquarter Hospital in Gilgit-Baltistan, Pakistan after ethical approval. The study duration was 6 months from August 2024 to January 2025. After taking informed consent, 227 children and young adolescents with ages ranging from 1 to 18 years were included using non-probability convenience sampling technique. Data was collected using a structured questionnaire including demographic factors, hygiene practices, healthcare access, and other relevant factors. Specimen bottles labeled with participant names and identification numbers were provided to parents and they were guided for stool sample collection. The collected specimens were processed in the Pathology Department of a healthcare facility for stool routine analysis. The data was analyzed using the Statistical Package for the Social Sciences (SPSS) version 25.

Results: Intestinal parasitic infections (IPIs) were present in 74(32.6%) children and young adolescents. *Ascaris lumbricoides* (46%) and *Giardia lamblia* (29.7%) were the most prevalent intestinal parasites. Access to clean drinking water and hygienic conditions had a significant relation with IPIs (p-value=0.001). No significant association was seen with other variables such as gender, age, education level, and family income.

Conclusion: Intestinal parasitic infections were prevalent in 32.6% children and young adolescents. *Ascaris lumbricoides* and *Giardia lamblia* were the most common parasites. There was a statistically significant influence of environmental factors, particularly access to clean drinking water and poor personal hygienic conditions, on the frequency of these infections.

Keywords: Parasitic infections. Ascariasis. Giardiasis.

INTRODUCTION

Intestinal parasitic infections pose a significant threat to global health, affecting 30% of the world population. Around 3.5 billion people have parasitic infestations across the world, out of which 450 million cases present with signs and symptoms. Approximately 200,000 deaths are attributed to these infections. The parasitic infections by hookworms cause 45000, *Ascaris* causes 4300, and *Entamoeba histolytica* causes 54000 annual deaths.^{1,2} Intestinal parasitic infections are responsible for 1.9 million disability-adjusted life years. The healthcare systems face significant financial constraints to cope with the rising number of IPIs.³

The major burden of IPIs lies in the developing countries.¹ The major reasons behind the greater prevalence of IPIs in these countries are poverty, overcrowding, insufficient handwashing practices, use of unwashed vegetables, lack of education &

awareness, poor hygiene, shortage of clean water supply, and inadequate healthcare facilities.⁴ Parasitic infections are caused by protozoans and helminths (worms). The main transmission route is fecal-oral, but transmission can also occur by larval penetration of skin.⁵ Children are prone to acquire IPIs owing to their weak immunity and poor hygienic practices. These infections cause abdominal pain, anemia, and malnutrition in children. They can also impair their physical and mental growth, leading to a great influence on productivity and quality of life. Children should be routinely screened for IPIs & their associated malnutrition and prescribed proper therapy, not only to reduce the morbidity & mortality attributed to IPIs but also to boost the physical and mental health of children.⁶

The prevalence of intestinal parasitic infections is very high in Pakistan, varying from 25% to 70%. Combating this high burden of IPIs is challenging owing to lack of awareness among the public, inadequate healthcare facilities, and a lack of a monitoring system for IPIs.⁷

The mountainous location of Gilgit-Baltistan with its harsh climate makes it a hard-to-reach area, leading to limited healthcare infrastructure in the region. The traditional practices of fetching water from far areas, lack of clean water availability, defecating in the open environment, and other unhygienic lifestyle conditions contribute to the favorable transmission

Sharif Medical & Dental College, Sharif Medical City,
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Uzma Ali
Senior Consultant Pathologist, Department of Pathology
Provincial Headquarter Hospital, Gilgit
E-mail: dr.uzma14@gmail.com

Received: September 5, 2025; Accepted: September 20, 2025

of parasites. In spite of the considerable risks involved, there is an absence of thorough data regarding the frequency and determinants of intestinal parasitic infections in Gilgit-Baltistan. The current study aims to address this critical gap by assessing the frequency of IPIs and associated factors in children and young adolescents in Gilgit-Baltistan to provide evidence for improving public health interventions, aligning with Pakistan's broader commitment to the Sustainable Development Goals.

METHODOLOGY

This descriptive, cross-sectional study was done on children and young adolescents visiting the Provincial Headquarter Hospital in Gilgit, Pakistan. The study duration was 6 months from August 2024 to January 2025. After ethical approval (Letter No. 1150/PHG/2023, 29-07-2024), 227 participants were included using non-probability convenience sampling technique. The sample size was estimated based on the expected prevalence rate of 30% IPIs, confidence level of 95% and margin of error of 6%.⁸ The inclusion criteria were children and young adolescents with an age range of 1 to 18 years, who were willing to participate and give stool samples. Children/adolescents on anti-parasitic/deworming medication and those with chronic gastrointestinal conditions unrelated to IPIs were excluded. Informed written consent was obtained from the parents or guardians before participation and they were explained the purpose of the study. Data was collected using a semi-structured questionnaire including demographic factors, education level of guardian, family income, access to clean drinking water, and physical hygiene conditions. Based on family income, those with monthly income <30,000 rupees were labeled as having low income, 30,000 to 70,000 rupees were middle income, and >70,000 rupees were high income. The physical hygienic conditions were categorized according to handwashing practices (before meals and after defecation), trimmed nails, regular wearing of shoes, in-home sanitation facility, eating raw fruits and vegetables, playing with soil, and swimming habit.⁹ Wide-mouthed, leak-proof specimen bottles labeled with participant names and identification numbers were provided to parents or guardians and they were guided for stool sample collection. Each participant was instructed to provide three stool samples on alternate days to increase the diagnostic sensitivity. The collected specimens were processed in the Pathology Department of a nearby healthcare facility for stool routine analysis by a trained laboratory

technologist. The fresh specimens received were physically examined and then two wet preparations were made using normal saline on one side and Lugol's iodine on the other side of the same glass slide. They were examined under 10X and 40X objective lens to identify ova, cysts or trophozoites of various parasites. In addition, wet preparations were also made by using the formalin-ether concentration technique. In this technique, 0.5 g of the stool sample was mixed with 10 ml of normal saline. The suspension was strained using a gauze, centrifuged, and the supernatant was discarded. Around 2.5 ml of 10% formaldehyde and 1 ml of ether were added to the sediment. The suspension was again centrifuged for 1-2 minutes at 3000 rpm. The supernatant was discarded and wet preparation was prepared from the sediment and examined under microscope.⁹ To ensure accuracy, each sample was examined independently by two trained microbiologists. In case of disagreement, the slides were re-examined together and senior microbiologists were also consulted for final confirmation. Positive cases were referred to the physician for treatment.

STATISTICAL ANALYSIS

The data was analyzed using the Statistical Package for the Social Sciences (SPSS) version 25. Categorical variables were presented as frequency & percentage, while numerical variables were summarized using mean and standard deviation. Association between categorical variables was determined using Pearson's Chi-square and Fisher's exact test. A p-value of <0.05 was considered statistically significant.

RESULTS

Out of 227 participants of our study, most (42.3%) were school-aged children (6-12 years), followed by young adolescents (33.5%), and children ≤5 years (24.2%). Most (55%) of the participants were males. A greater proportion of the participants (41.4%) had low income, 36.6% had middle income, and 22% had high income. Regarding the education status of guardians, 42.8% were illiterate, 34.8% were educated till secondary, and 22.4% had completed graduation. The majority (56.4%) of the population had no access to clean water and reported good physical hygienic conditions (62.1%).

Out of 227 participants, 74(32.6%) had parasitic infestations while 153(67.4%) showed no infestation. Access to clean drinking water and hygienic conditions were significantly associated with infestation rates (p-value=0.001). A higher

frequency of IPIs was observed among participants with no access to clean drinking water and poor physical hygienic conditions. No significant association was seen with other variables such as gender, age, education level, and family income (Table 1).

Regular swimming habits, playing with soil, and eating raw fruits & vegetables were significantly associated with the presence of parasitic infestations

($p=0.001$). More frequent IPIs were exhibited by those who did not trim nails, wear shoes or wash hands before meals, and after defecation regularly ($p=0.001$).

The most frequently reported parasite was *Ascaris lumbricoides* (46%), followed by *Giardia lamblia* (29.7%). Other parasites, such as *Hymenolepis nana*, *Taenia saginata*, *Trichuris trichiura*, and *Entamoeba histolytica*, had lower frequency rates (Figure 1).

Table 1: Association of Socio-Demographic and Environmental Factors with Parasitic Infestation

Socio-Demographic and Environmental Factors		Parasitic Infestation (Frequency & Percentage)			p-value
		Present (n=74)	Absent (n=153)	Total	
Gender	Male	38(16.7%)	87(38.3%)	125(55%)	0.43
	Female	36(15.9%)	66(29.1%)	102(45%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	
Age Groups (Years)	≤5	23(10.1%)	32(14.1%)	55(24.2%)	0.18
	6 to 12	26(11.5%)	70(30.8%)	96(42.3%)	
	13 to 18	25(11%)	51(22.5%)	76(33.5%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	
Education Level of Guardian	Illiterate	29(12.8%)	68(30%)	97(42.8%)	0.66
	Secondary	26(11.5%)	53(23.3%)	79(34.8%)	
	Graduation	19(8.3%)	32(14.1%)	51(22.4%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	
Family Income	Low	29(12.8%)	65(28.6%)	94(41.4%)	0.89
	Middle	28(12.3%)	55(24.2%)	83(36.5%)	
	High	17(7.5%)	33(14.5%)	50(22%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	
Access to Clean Drinking Water	No	65(28.6%)	63(27.8%)	128(56.4%)	0.001*
	Yes	9(4%)	90(39.6%)	99(43.6%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	
Physical Hygienic Conditions	Poor	58(25.6%)	28(12.3%)	86(37.9%)	0.001*
	Good	16(7%)	125(55.1%)	141(62.1%)	
	Total	74(32.6%)	153(67.4%)	227(100%)	

*Significant p-value




Figure 1: Species Distribution of Intestinal Parasitic Infections

DISCUSSION

Intestinal parasitic infections remain a critical public health challenge globally, especially in low-resource settings where factors such as inadequate sanitation, unsafe water, and poor hygiene practices are widespread. The prevalence of IPIs had also increased due to food globalization and frequent travel to various countries.^{10,11} The findings of our study revealed a significant burden of IPIs (32.6%) among children and young adolescents in Gilgit-Baltistan, Pakistan. In two studies done in Ethiopia, IPIs were present in 24.4% of the children.^{12,13} Other studies reported 28.27% and 29.4% prevalence of parasitic infestation among children in Ethiopia.^{14,15} Sitotaw et al. reported a higher (62.40%) rate of parasitic infections among school children in Ethiopia.¹⁶ This showed that the prevalence of IPIs can vary greatly from one area to another, even within the same country. In Argentina, 55.4% of the participants aged 1-15 years had IPIs.¹⁷ In Somalia, IPIs were reported in 82.9% of the children.¹⁸ A study from Lower Dir, Pakistan showed an 82% IPI rate among children. The high frequency of IPIs in children and young adolescents was attributed to their increased vulnerability due to unhygienic behaviors and greater interaction with contaminated environment.¹⁹ On the other hand, a lower prevalence of IPIs had been found in Europe (5.9%).²⁰ In a study in India, the frequency of IPIs was 9.3%.²¹

The most common parasites in our study were *Ascaris lumbricoides* (46%) and *Giardia lamblia* (29.7%). Other parasites isolated were *Hymenolepis Nana* (8.1%), *Taenia saginata* (6.7%), *Trichuris trichiura* (4.1%), and *Entamoeba histolytica* (5.4%). In Lower Dir from Pakistan, *Ascaris* was the most common parasite isolated (57.7%), followed by hookworm (40.7%), *Taenia saginata* (20.9%), *Enterobius* (14.8%), *Trichuris* (14.5%), *Hymenolepis* (12.9%), and *Entamoeba* (9.25%).¹⁹ Abebaw et al. and Scavuzzo et al. reported *Giardia lamblia* and *Hymenolepis nana* as the most common causes of parasitic infections.^{12,17} In Ethiopia, parasitic infestations were mainly caused by *Entamoeba histolytica* (10.9%) and *Schistosoma mansoni* (7.4%).¹³ According to a study by Duguma et al., *Ascaris* was the most common parasite (8%), followed by *Trichuris* (6.2%) and *Giardia* (4%).¹⁵ In another study, the most common parasites were *Ascaris* (22.7%), Hookworms (20.6%), and *Entamoeba histolytica* (8.1%). Other parasites isolated were *Trichuris trichiura* (7.6%), *Giardia* (6.5%), *Hymenolepis nana* (5.7%), and *Schistosoma*

mansoni (4.4%).¹⁶ In Somalia, the most common parasites were *Ascaris* (46.6%), *Giardia lamblia* (22.1%), and *Entamoeba histolytica* (17.6%).¹⁸ In India, *Ascaris* (57%) and hookworm (42%) were the most common parasites.²¹

In our study, males comprised 55% of the population, and 45% were females. There was no statistically significant difference in parasitic infestation rates between males and females. Similarly, there were 53.2% males and 46.8% females in a study done by Yeshitila et al. Females were predominantly infected but with no statistical difference.¹³ Similarly, another study revealed that IPIs affected males and females equally.¹⁶ Our study showed that the frequency of IPIs was almost similar in various age groups and the age factor did not significantly affect infestation rates ($p=0.18$). Our results were in accordance with a study by Sitotaw et al., where no statistically significant difference in IPIs in different age groups was observed.¹⁶ The IPIs affected a greater proportion of participants in the 10-14 years age groups (26.7%) in a study by Yeshitila et al.¹³ In another study, there was a significant difference in the frequency of IPIs in different age groups, with <5 years most commonly affected, followed by 6-12 years.¹⁷ Osman et al. from Somalia reported that IPIs were significantly prevalent in 13-36 months age group.¹⁸ The results of the current study reported no significant association between the infestation rates and the education level of guardians ($p=0.66$). Similarly, family income did not significantly impact infestation rates ($p=0.89$). In contrast, a study revealed a significant association of IPIs with family income and education level. The rate of IPIs was significantly higher in the low income group and those with lower literacy.¹⁶

In our study, the factors significantly associated with IPIs were limited access to clean drinking water and poor physical hygienic conditions. Children without access to clean water (28.6%) had a significantly higher infection rate as compared to those with access to clean water (4%) ($p=0.000$). Poor hygienic conditions were significantly associated with higher infection rates (25.6%) as compared to only 7% rate among those with good hygienic practices. ($p=0.000$). Duguma et al. showed that access to clean drinking water, lack of handwashing, waste disposal methods, habit of wearing shoes, and poor hygiene were significantly associated with IPIs in Ethiopia.¹⁵ In other studies, the most common risk factors associated with parasitic infections in children were untrimmed fingernails and lack of

handwashing.^{12,14} The predisposing factors for parasitic infestation were lack of handwashing and swimming in contaminated water in a study.¹³ Scavuzzo et al. reported that overcrowding and walking barefoot, were significantly associated with IPIs in Argentina.¹⁷

CONCLUSION

Intestinal parasitic infections were prevalent in 32.6% children and young adolescents. *Ascaris lumbricoides* and *Giardia lamblia* were the most common parasites. There was a statistically significant influence of environmental factors, particularly access to clean drinking water and poor hygienic conditions on the frequency of these infections.

LIMITATIONS & RECOMMENDATIONS

Our study had a few limitations, such as cross-sectional design, a single-centered study, and the use of convenience sampling, limiting the generalizability of the findings. The reliance on self-reported data for hygiene practices introduced the possibility of recall bias or social desirability bias. The study was conducted over 6 months, but parasitic prevalence may vary seasonally, which was not discussed.

The study recommended that targeted public health interventions focusing on improving water & sanitation infrastructure and promoting hygiene education should be implemented. Routine deworming programs, particularly in schools, are essential for reducing the burden of parasites such as *Ascaris lumbricoides*, etc. Health education campaigns focused on promoting hygiene practices, such as handwashing and safe food handling, can further reduce the spread of parasitic infections.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

U.A: Performed data analysis, interpretation, and drafted the initial manuscript.

S.N: Contributed to study design, methodology development, and critical revision of the manuscript.

A.Q: Conceived and designed the study, supervised data collection.

A.A: Participated in data collection, laboratory work, and tabulation of results.

N.Y: Assisted in literature review, data entry, and formatting of figures and tables.

Z.W: Supervised the overall project, reviewed the final manuscript critically for important intellectual content, and approved it for publication.

REFERENCES

1. Elmonir W, Elaadli H, Amer A, El-Sharkawy H, Bessat M, Mahmoud SF, et al. Prevalence of intestinal parasitic infections and their associated risk factors among preschool and school children in Egypt. *PLoS One*. 2021; 16(9):e0258037. doi:10.1371/journal.pone.0258037.
2. Hajare ST, Gobena RK, Chauhan NM, Erniso F. Prevalence of intestinal parasite infections and their associated factors among food handlers working in selected catering establishments from Bule Hora, Ethiopia. *Biomed Res Int*. 2021; 2021:6669742. doi:10.1155/2021/6669742.
3. Gebru H, Deyissia N, Medhin G, Kloos H. The association of sanitation and hygiene practices with intestinal parasitic infections among under-14 children in rural Dire Dawa, Eastern Ethiopia: a community based cross-sectional study. *Environ Health Insights*. 2023; 17:11786302231180801. doi:10.1177/11786302231180801.
4. Getie M, Belay G, Amare A, Abebe W, Deress T. Burden and factors influencing intestinal parasitic infections among food handlers in Gondar City, Northwest Ethiopia. *Front Public Health*. 2024; 12:1362086. doi:10.3389/fpubh.2024.1362086.
5. Mussema A, Tafesse W, Temam L. Prevalence of intestinal parasite infection among prison inmates and their associated risk factors at Hosanna Town, South-Central Ethiopia: a cross-sectional study. *J Parasitol R*. 2024; 2024:1-8. doi:10.1155/2024/767774.
6. Bisetegn H, Debash H, Ebrahim H, Erkihun Y, Tilahun M, Feleke DG. Prevalence and determinant factors of intestinal parasitic infections and undernutrition among primary school children in North-Central Ethiopia: a school-based cross-sectional study. *J Parasitol Res*. 2023; 2023:2256910. doi:10.1155/2023/2256910.
7. Fauziah N, Aviani JK, Agrianfanny YN, Fatimah SN. Intestinal parasitic infection and nutritional status in children under five years old: a systematic review. *Trop Med Infect Dis*. 2022; 7(11):371. doi:10.3390/tropicalmed7110371.
8. Irum S, Ahsan A, Ahmed H, Khan A, Yayi G, Mehboob M, et al. A demographic survey on the prevalence of gastrointestinal parasites based on socioeconomic determinants in Pakistan. *J Infect Dev Ctries*. 2021; 15(11):1738-43. doi:10.3855/jidc.12032.
9. Birhanu AA, Yimer M, Debash H, Abate MA. Prevalence and associated factors of intestinal parasitic infections and undernutrition among elementary school children in Zenzelima town, Northwest Ethiopia: a school-based cross-sectional study. *Health Sci Rep*. 2025; 8(7):e70964. doi:10.1002/hsr2.70964.

10. Shan F, Khan A, Khan H, Ali H, Khan MK, Rahat A. Detecting intestinal parasites in diarrheal cases among children in Dir Lower District. *JHRR*. 2024; 4(1):1281-5. doi:10.61919/jhrr.v4i1.537.
11. Ahmed M. Intestinal parasitic infections in 2023. *Gastroenterology Res*. 2023; 16(3):127-40. doi:10.14740/gr1622.
12. Abebaw A, Alemu G, Ayehu A. Prevalence of intestinal parasites and associated factors among children from child centres in Bahir Dar city, Northwest Ethiopia. *Trop Doct*. 2020; 50(3):194-8. doi:10.1177/0049475520920161.
13. Yeshitila YG, Zewde H, Mekene T, Manilal A, Lakew S, Teshome A. Prevalence and associated risk factors of intestinal parasites among school-children from two primary schools in Rama Town, Northern Ethiopia. *Can J Infect Dis Med Microbiol*. 2020; 2020(1):5750891. doi:10.1155/2020/5750891.
14. Girma A, Genet A. Magnitude and determinants of intestinal parasites among children under five in Ethiopia during 2010-2023: a systematic review and meta-analysis. *Fetal Pediatr Pathol*. 2024; 43(1), 47-65. doi:10.1080/15513815.2023.2284920.
15. Duguma T, Worku T, Sahile S, Asmelash D. Prevalence and associated risk factors of intestinal parasites among children under five years of age attended at Bachuma Primary Hospital, West Omo Zone, South-west Ethiopia: a cross-sectional study. *J Trop Med*. 2023; 2023:2268554. doi:10.1155/2023/2268554.
16. Sitotaw B, Shiferaw W. Prevalence of intestinal parasitic infections and associated risk factors among the first-cycle primary schoolchildren in Sasiga District, Southwest Ethiopia. *J Parasitol Res*. 2020; 2020:8681247. doi:10.1155/2020/8681247.
17. Scavuzzo CM, Campero MN, Oberto MG, Porcasi X, Periago MV. Intestinal parasites in children from native communities of Salta, Argentina. *Colomb Med (Cali)*. 2024; 55(1):e2025948. doi:10.25100/cm.v55i1.5948.
18. Osman MM, Hussain SA, Omar MA, Isaack MM, Hanafi AMH, Ahmed MY, et al. Pattern of intestinal parasitic infections in children with malnutrition in Somalia. *IJID Regions*. 2024; 13: 100431. doi:10.1016/j.ijregi.2024.100431.
19. Ulhaq Z, Khan W, Khan MF, Kabir M, Ujjan AA, Ullah W, et al. Prevalence of intestinal parasitic diseases in school children of rural areas of district Lower Dir, Pakistan. *Braz J Biol*. 2021; 82:e243150. doi:10.1590/1519-6984.243150.
20. Kantzanou M, Karalexi MA, Vrioni G, Tsakris A. Prevalence of intestinal parasitic infections among children in Europe over the last five years. *Trop Med Infect Dis*. 2021; 6(3):160. doi:10.3390/tropicalmed6030160.
21. Ulaganeethi R, Rajkumari N, Gururajan A, Gunalan A, Langbang D, Kumar G. Intestinal parasitic infections and its trends: a 5-year findings from a tertiary care centre, Puducherry, South India. *J Parasit Dis*. 2021; 45(2):400-5. doi:10.1007/s12639-020-01310-9.



Diagnostic Accuracy of Retrograde Urethrogram in Determination of Urethral Stricture taking Cystoscopy as Gold Standard

Rafia Irum, Aneez Qayyum, Hira Bushra, Hamna Mazhar

ABSTRACT

Objective: To assess the diagnostic accuracy of retrograde urethrogram in determination of urethral strictures taking cystoscopy as the gold standard.

Methodology: This cross-sectional study was conducted at the Department of Radiology, Sharif Medical & Dental College, Lahore from May to October 2024. After ethical approval and written informed consent, 189 patients meeting the inclusion criteria were recruited through non-probability convenience sampling technique. These patients underwent retrograde urethrogram (RUG) with contrast injection, followed by cystoscopy 4-5 days later. Demographic details and history of risk factors were recorded. Radiographic findings were assessed by a radiologist, and patients were classified as positive or negative for urethral strictures. Data were analyzed using Statistical Package for the Social Sciences (SPSS) version 25. A 2x2 table was made and diagnostic accuracy was calculated. Chi-square test was used to compare categorical variables.

Results: Retrograde urethrogram detected urethral strictures in 24.3% of cases with a sensitivity of 75.56%, specificity of 91.67%, positive predictive value (PPV) of 73.91%, and negative predictive value (NPV) of 92.31%. The overall diagnostic accuracy of RUG was 87.83%. Age groups, history of urinary tract infections, and prostate surgeries had significant association with the presence/absence of strictures on RUG ($p < 0.05$).

Conclusion: The diagnostic accuracy of retrograde urethrogram was high in determination of urethral strictures taking cystoscopy as the gold standard. Urethral strictures were significantly more prevalent among older males with history of urinary tract infections and prostate surgeries.

Keywords: Urethral stricture. Cystoscopy. Contrast media. Iohexol.

INTRODUCTION

The male urethra is an essential part of the urinary system, measuring approximately 20 cm in length, and it is surrounded by the corpus spongiosum, a vascular tissue that plays a critical role in maintaining the structure and function of the urethra. The urethra itself is divided into two primary sections: the anterior urethra and the posterior urethra, which are separated by the membranous urethra.¹ Each part of the urethra serves a unique function in the passage of urine from the bladder to the outside of the body.²

Urethral stricture is a condition characterized by the narrowing of the urethra, which can have a significant impact on a patient's quality of life. This condition develops slowly over time and can result in a variety of symptoms, including dysuria, weak urinary stream, incomplete bladder emptying, and urinary tract infections (UTIs).³ In recent years, the incidence of iatrogenic urethral strictures has been

increasing, primarily due to the rise in minimally invasive, transurethral urological procedures like cystoscopy and transurethral resection of the prostate (TURP).⁴ Chronic inflammatory conditions such as lichen sclerosis, urinary tract infections, prolonged catheterization, and the increased cases of urethral traumas resulting from traffic accidents and workplace injuries are among other contributing factors.⁵ Many complications including acute urinary retention, prostatitis, epididymo-orchitis, hydronephrosis, chronic kidney disease, peri-urethral abscess, and kidney stones can arise if the disease remains untreated for a longer duration.⁶

Various diagnostic techniques like uroflowmetry, cystoscopy, RUG, voiding cysto-urethrography (VCUG) can be useful in making diagnosis of urethral strictures and describing key characteristics like position & length of stricture and severity of condition. Cystoscopy, usually performed under anesthesia, makes the definitive diagnosis of urethral stricture. However, it cannot assess the length of the stricture and does not provide information about the condition of the surrounding tissue.⁴ Retrograde urethrogram technique offers several advantages including its ability to provide real-time imaging of the urethra as contrast material is injected, allowing for dynamic assessment of the stricture's characteristics. It also enables the clinician to measure its length, which is a crucial factor in

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Rafia Irum
Associate Professor, Department of Radiology
Sharif Medical & Dental College, Lahore
E-mail: dr.rafiassim@gmail.com

Received: July 10, 2025; Accepted: September 1, 2025

planning further treatment, such as urethral dilation, urethrotomy, or urethroplasty.⁷

A recent high number of urethral strictures is being reported in Pakistan due to increased prevalence of sexually transmitted diseases and improper catheterization techniques. Retrograde urethrogram is comparatively inexpensive, and less invasive technique for its diagnosis. It was pertinent to compare the diagnostic accuracy of RUG against the gold standard (cystoscopy) in our local setting to develop locally relevant diagnostic guidelines and improve patient outcomes.

METHODOLOGY

After approval from the institutional review board (Letter No. SMDC/SMRC/337-24, 22-05-2024), this cross-sectional study was carried out at the Department of Radiology, Sharif Medical & Dental College, Lahore from May to October 2024. A sample size of 189 was estimated using 90% confidence level, 7% margin of error, 100% sensitivity of RUG, 66.7% specificity, and prevalence of urethral strictures at 28.5%.⁸ The male patients aged 18-75 years, who presented with restricted urinary flow from the bladder, urinary tract infection, or inflammation of the bladder, incomplete voiding, and/or painful urination were enrolled by using non-probability convenience sampling technique. Patients who had suggestive symptoms of acute urethritis, malignancy or metastatic disease, and received renal implant were excluded.

After obtaining written informed consent, demographic and risk factors detail [name, age, body mass index (BMI), duration of symptoms, marital status, occupation, history of diabetes, hypertension, smoking, urinary tract infection, and previous prostate surgery] were recorded on a questionnaire. The patients then underwent retrograde urethrogram in the right oblique or semi-lateral position after emptying the bladder. The right leg was partially flexed, with the left leg stretched above it and the left hip just above the table. A Foley's catheter was placed in the external meatus after disinfecting the glans, and 10-15 ml of 15% (w/v) Iohexol was injected through catheter. The patients were told to relax their pelvic floor muscles during the injection. The penile tip was pressed tightly around the catheter during injection to prevent the escape of contrast medium. A single spot film was taken with a constant film focus distance of 1 meter maintained for all radiographs. All scans were performed and interpreted by two consultant radiologists with at least 4 years of experience in radiology to improve accuracy. Patients were labeled as positive for

urethral strictures if annular or longitudinal narrowing of the urethra was seen on radiographs.⁹ After 4-5 days, cystoscopy was performed through the suprapubic tract, and the cystoscope was advanced through the bladder neck to the level of the stricture. Findings were recorded, and patients were confirmed for presence or absence of urethral strictures. The cases that were positive on both RUG and cystoscopy were labelled as true positives (TP). Those who were positive on RUG but negative on cystoscopy were false positives (FP). False negatives (FN) were negative on RUG but positive on cystoscopy. Those who were negative on both the tests were true negatives (TN).

STATISTICAL ANALYSIS

Data was entered and analyzed using Statistical Package for the Social Sciences (SPSS) version 25. Descriptive statistics like mean±standard deviation and frequency (percentage) were used for numeric and categorical variables, respectively. A 2x2 table was made by taking cystoscopy as a gold standard. The sensitivity, specificity, PPV, NPV, and diagnostic accuracy of RUG were calculated. The categorical variables were compared using Chi-square test. A p-value of <0.05 was considered statistically significant.

RESULTS

The study included 189 male participants and the mean age of the participants was 46.87±16.74 years, indicating a wide range of ages (from 30.9 to 63.6 years). Most of the participants (56.6%) belonged to 18-50 years age group. Nearly half (48.1%) of them were married and only 45% were employed. The mean BMI was 27.31±4.87 kg/m², suggesting that the majority of the participants were in the overweight range. The mean duration of symptoms was 11.70±6.72 weeks, reflecting that some participants experienced longer duration of symptoms than others. The frequency of urethral stricture on RUG was 24.3%. The age groups showed significant association with urethral strictures on RUG. Most of the patients with urethral strictures were older than 50 years (Table 1).

The risk factors profile of 189 participants revealed 47.6% diabetics, 40.7% hypertensive and 49.7% smokers. None of these factors had any significant association with urethral strictures on RUG ($p \geq 0.05$). Urethral stricture was significantly more prevalent among those with UTIs (16.9%) and history of prostate surgery (15.9%) (Table 1).

The cross-tabulation of cystoscopy and RUG outcomes showed significant results (Table 2).

Retrograde urethrogram showed a sensitivity of 75.56%, specificity of 91.67%, PPV of 73.91%, NPV of 92.31%, and an overall diagnostic accuracy of 87.83%.

DISCUSSION

This study aimed to evaluate the diagnostic accuracy of retrograde urethrography for detecting urethral strictures keeping cystoscopy as gold standard. The frequency of urethral stricture on RUG was 24.3%. The mean age of the participants was 46.87 ± 16.74 years and most of the patients with urethral strictures on RUG were older than 50 years age ($p=0.00001$). Similarly, de Farias RB et al. showed that urethral strictures were significantly more frequent (82.2%) among patients over 40 years of age ($p < 0.001$).¹⁰ However, a previous study didn't reported any significant association of age with the occurrence of urethral strictures among post-TURP patients ($p=0.932$).¹¹ The results of the current study showed that hypertension, diabetes mellitus, and smoking had no significant association with the occurrence of urethral stricture on RUG. Contrary to our findings, diabetes mellitus ($p=0.038$) and hypertension ($p=0.012$) were significant risk factors in recurrence

of urethral strictures among patients who underwent internal urethrotomy.¹²

Our significant results also revealed that the majority of cases of urethral strictures diagnosed on RUG had a history of UTI and prostate surgeries. Another study also reported that TURP represented 41% of all iatrogenic causes of urethral strictures. Infections accounted for 26.6% of cases.¹³ Sekar et al. found that 12.8% patients who underwent TURP developed urethral strictures. They also revealed that the use of larger size Foley's catheter during this procedure was significantly associated with post-operative urethral strictures.¹⁴

When compared with cystoscopy as gold standard in our study, RUG had a sensitivity of 75.56%, specificity of 91.67%, positive predictive value of 73.91%, negative predictive value of 92.31%, and an overall accuracy of 87.83%. In a previous study, when RUG was compared with intraoperative findings as gold standard, it showed 100% sensitivity, 66.7% specificity and an overall accuracy of 83.3% in detecting urethral strictures.⁸ Patel et al. also documented in their review of literature that RUG was the best technique in making diagnosis of urethral structures following urethral traumas.¹⁵

Table 1: Association of Risk Factors with Urethral Strictures on RUG (n=189)

Risk Factors		Urethral Stricture on RUG		Total	p-value
		Present (n=46)	Absent (n=143)		
Age Groups (Years)	16-50	09(4.8%)	98(51.9%)	107(56.7%)	0.00001*
	51-70	37(19.5%)	45(23.8%)	82(43.3%)	
	Total	46(24.3%)	143(75.7%)	189(100%)	
Urinary Tract Infections	Yes	32(16.9%)	57(30.2%)	89(47.1%)	0.0004*
	No	14(7.4%)	86(45.5%)	100(52.9%)	
	Total	46(24.3%)	143 (75.7%)	189(100%)	
History of Prostate Surgery	Yes	30(15.9%)	58(30.7%)	88(46.6%)	0.003*
	No	16(8.4%)	85(45%)	101(53.4%)	
	Total	46(24.3%)	143(75.7%)	189(100%)	

*Significant p-value

Table 2: Cross Tabulation for Determining Diagnostic Accuracy of RUG in Diagnosis of Urethral Strictures taking Cystoscopy as Gold Standard

Retrograde Urethrogram	Cystoscopy (Gold Standard)		Total	p-value
	Positive	Negative		
Positive	34(18%) (TP)	12(6.3%) (FP)	46(24.3%)	0.00001*
Negative	11(5.8%) (FN)	132(69.9%) (TN)	143(75.7%)	
Total	45(23.8%)	144(76.2%)	189(100%)	

*Significant p-value

In contrast to our findings, a study reported that RUG was used in combination with VCUG for the evaluation of urethral strictures. They found that combined preoperative use of RUG and VCUG underestimated urethral stricture measurements, especially in the membranous and bulbar regions, when compared with intraoperative results ($p < 0.05$).¹⁶ A high level of diagnostic accuracy is especially important as the precise location and length of the stricture are critical for planning surgical interventions like simple urethral dilation, an internal urethrotomy, or a more complex urethral reconstruction, i.e. urethroplasty. Misjudging the length or location of the stricture can lead to suboptimal outcomes, such as incomplete stricture excision or recurrence of the stricture postoperatively.^{16,17}

Despite its high diagnostic accuracy reported in the literature, RUG had limitations, particularly in its ability to provide critical and detailed information regarding the fibrosis of soft tissues surrounding the stricture, such as spongiofibrosis or periurethral fibrosis. This condition required a more extensive reconstruction, and failing to detect it preoperatively could result in inadequate treatment.¹⁸ To address this limitation, alternative imaging modalities such as sonourethrography (SUG) and magnetic resonance urethrography (MRU) have been explored. Bogdanov et al. found that stricture measurements on RUG were significantly different than those of intraoperative findings. However, MRU results were significantly similar to those of intraoperative findings.¹⁹ A study documented that MRU was significantly most accurate in preoperative evaluation of strictures, followed by cystourethrography (RUG and VCUG) when compared to intraoperative measurements. The least accurate was SUG in this regard.⁹ In contrast, a previous study showed that SUG was the most accurate method with a sensitivity of 60% and specificity 87.8% when compared with VCUG and cystoscopy for determining the length of strictures.¹⁷ Jesrani et al. also highlighted that SUG had a higher diagnostic accuracy in determination of urethral strictures as compared to RUG.²⁰ With advancements in technologies, artificial intelligence and machine learning were more accurate in the detection and reconstruction of urethral strictures.²¹ Our findings reflected the broad utility of RUG in clinical practice, where it can be used for assessing urethral strictures providing crucial information about the location, length, and severity of the stricture.

CONCLUSION

When compared with cystoscopy as gold standard, retrograde urethrography had a good diagnostic accuracy in determination of urethral strictures among males. Urethral strictures were significantly more prevalent among older males with history of urinary tract infections and prostate surgeries.

LIMITATIONS & RECOMMENDATIONS

This study had a few limitations: cross-sectional design, single-centered study, non-probability convenience sampling technique limiting the generalizability of results. This study also lacked information on detailed characteristics of the stricture and post-operative follow-up. Further research is needed to explore the role of advanced imaging modalities and the integration of RUG with other techniques in the comprehensive evaluation of urethral strictures, particularly in complex cases where precise assessment of soft tissue involvement is necessary.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

R.I: Supervised the overall project, reviewed the final manuscript critically for important intellectual content, and approved it for publication.

A.Q: Contributed to study design, methodology development, and critical revision of the manuscript.

H.B: Performed data analysis, interpretation, and drafted the initial manuscript.

H.M: Conceived and designed the study, supervised data collection.

REFERENCES

1. Hassan U, Bacha R, Khan MF, Laique T. Diagnostic accuracy of sonourethrography for anterior urethral stricture taking retrograde urethrogram as gold standard. *PJMHS*. 2020; 14(4):1861-3. Available from: <http://pjmhs online.com/published-issues/2020-issues/oct-dec-2020/20-12-1861>.
2. Surgical anatomy of the penis and urethra. In: Hadidi AT, editor. *Hypospadias surgery: an illustrated textbook*. 2nd Edition. Springer International Publishing. 2022:105-26. doi:10.1007/978-3-030-94248-9_4.
3. Yan Y, Wu Y, Li A, Yang A, Tao J, Wang X. Impact of a clinical care pathway developed through the action research method on the psychological well-being and quality of life in male patients with urethral stricture. *Medicine (Baltimore)*. 2024; 103(9):e37321. doi:10.1097/MD.00000000037321.

4. Mikolaj F, Karolina M, Oliwia K, Jakub K, Adam K, Mariusz B, et al. Retrograde urethrography, sonourethrography and magnetic resonance urethrography in evaluation of male urethral strictures. Should the novel methods become the new standard in radiological diagnosis of urethral stricture disease? *Int Urol Nephrol*. 2021; 53(12):2423-35. doi:10.1007/s11255-021-02994-5.
5. Cavalcanti A, Schul A, Restrepo CF, Aldred JE. Pathophysiology of urethral stricture: optimizing urethroplasty outcomes. In: Hofer MD, editor. *Scientific Advances in Reconstructive Urology and Tissue Engineering*; Academic Press. 2022; 177-95. doi:10.1016/B978-0-323-91199-3.00010-4.
6. Basaif WS, Alamri HH, Mousa HW, Alsayed RA, Almohammadi AM, Altulayqi WH, et al. Types of urethral stricture and their recurrence rates post urological treatments. *Int J Community Med Public Health*. 2021; 8(7):3634-7. doi:10.18203/2394-6040.ijcmph20212347.
7. Harris D, Zhou C, Girardot J, Kidron A, Gupta S, Cavalcanti AG, et al. Imaging in urethral stricture disease: an educational review of current techniques with a focus on MRI. *Abdom Radiol (NY)*. 2023; 48(3):1062-78. doi:10.1007/s00261-022-03761-5.
8. Kelkar TS, Joshi A. Comparing retrograde urethrography readings with intraoperative findings in urethral stricture. *Int J Res Med Sci*. 2019; 7(6):2333-6. doi:10.18203/2320-6012.ijrms20192523.
9. Freitas PS, Alves AS, Correia PS, Dias JL. Urethrocytography: a guide for urological surgery? *Diagn Interv Radiol*. 2023; 29(1):9-17. doi:10.5152/dir.2022.21640.
10. de Farias RB, Neto FT, de Aguiar Cavalcanti G, Martins FE, Lima SV. Evaluation of the etiological profile, age and findings in retrograde and voiding urethrocytography of men with urethral stricture. *Scientific Reports*. 2025; 15(1):5935. doi: 10.1038/s41598-025-89389-z.
11. Durmus E, Ok F, Sert IU. The factors affecting occurrence of urethral stricture after transurethral resection of the prostate. *J Clin Med Kaz*. 2022; 19(6):68-72. doi:10.23950/jcmk/12690.
12. Gul A, Ekici O, Zengin S, Barali D, Keskin T. Investigation of risk factors in the development of recurrent urethral stricture after internal urethrotomy. *World J Clin Cases*. 2024; 12(14):2324-31. doi:10.12998/wjcc.v12.i14.2324.
13. Abdeen BM, Leslie SW, Badreldin AM. Urethral strictures. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing. 2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK564297>.
14. Sekar H, Palaniyandi V, Krishnamoorthy S, Kumaresan N. Post-transurethral resection of prostate urethral strictures: are they often underreported? A single-center retrospective observational cohort study. *Urol Ann*. 2021; 13(4):329-35. doi:10.4103/UA.UA_165_19.
15. Patel AB, Osterberg EC, Satarasinghe PN, Wenzel JL, Akbani ST, Sahi SL, et al. Urethral injuries: diagnostic and management strategies for critical care and trauma clinicians. *J Clin Med*. 2023; 12(4):1495. doi:10.3390/jcm12041495.
16. Salimi H, Menbari Oskouie I, Mohammadi R, Nazarpour MJ, Niknam N, Nikoubakht MR, et al. Retrograde urethrography (RUG) combined with voiding cystourethrography (VCUG) versus surgical findings in assessment of urethral strictures length. *Urologia*. 2025; 92(2):342-7. doi:10.1177/03915603241292840.
17. Palau Roig M, Botto Lugo SM, Martinez Albaladejo P, Utiel Atienzar A, Cuenca Ramirez MLD, Gonzalez Diez D, et al. Comparative utility of urethrosonography versus classic techniques in diagnosing urethral stricture: a single center study. *Abdom Radiol (NY)*. 2025. doi: 10.1007/s00261-025-05054-z.
18. Frankiewicz M, Vetterlein MW, Markiet K, Adamowicz J, Campos-Juanatey F, Cocci A, et al. Ultrasound imaging of male urethral stricture disease: a narrative review of the available evidence, focusing on selected prospective studies. *World J Urol*. 2024; 42(1):32. doi:10.1007/s00345-023-04760-x.
19. Bogdanov AB, Katibov MI, Goncharuk DA, Veliev EI, Sokolov AE, Ivkin EV, et al. IP22-08 safety and capabilities of short static MR-urethrography for assessment of urethral strictures length and spongiositis grade. *J Urol*. 2025; 213(5S):e1170. doi:10.1097/01.JU.0001110136.41716.b7.08.
20. Jesrani A, Faiq SM, Hussain M, Lalwani A, Farid M, Saad M. Comparison of sonourethrography and retrograde urethrography in anterior urethral strictures taking surgical findings as gold standard. *PJR*. 2020; 30(3):158-63. Available from: <https://www.pakjr.com/index.php/PJR/article/view/1227/826>.
21. Wong HPN, So WZ, Fong KY, Tiong HY, Kulkarni S, Castellani D, et al. Advances in urethral stricture diagnostics and urethral reconstruction beyond traditional imaging: a scoping review. *Cent European J Urol*. 2024; 77(3):528-37. doi:10.5173/ceju.2024.121.



Comparison of Effectiveness between Diclofenac Suppositories and Injection Diclofenac in Abdominal Surgeries

Muhammad Abu Usama, Fizza Laraib, Farhana Manzoor, Maham Fatima, Syed Suleman Zafar Jafary, Hafiz Zain-ul-Abdeen

ABSTRACT

Objective: To compare the effectiveness of diclofenac suppositories and injection diclofenac in pain relief after abdominal surgeries.

Methodology: This quasi-experimental study was conducted at the West Surgical Ward, Mayo Hospital, Lahore from June 2022 to January 2023 after taking approval from ethical committee. Total 136 patients undergoing elective and emergency abdominal surgeries were enrolled in the study via non-probability convenience sampling technique. Informed written consent was taken from all the patients. The patients were divided into two groups (group A & B). Group A was given diclofenac suppositories and group B was given injection diclofenac via the intramuscular (I/M) route. The pain scores were recorded by using visual analogue scale (VAS) at 0, 8, 16, and 24 hours. The data was analyzed using Statistical Package for the Social Sciences (SPSS) version 26.

Results: There was a significant decrease in mean pain scores at 0, 8, 16, and 24 hours post-operatively in group A as compared to group B. The maximum difference in the means was observed at 16 hours post-operatively ($p < 0.001$). Only 15(11.1%) patients in group A required rescue analgesia, while more than half the patients in group B required rescue analgesia for effective pain relief ($p = 0.001$). The proportion of patients experiencing adverse effects of the drug in group A was significantly smaller (8.1%) as compared to group B (25.7%) ($p = 0.001$).

Conclusion: Rectal suppositories of diclofenac were significantly more effective for post-operative pain relief among patients undergoing abdominal surgeries as compared to diclofenac intramuscular injection.

Keywords: Pain. Post-operative. Visual Analog Scale. Analgesia. Diclofenac.

INTRODUCTION

Abdominal surgeries have a major role in the treatment of cases of non-traumatic acute abdominal pain. Appendicitis is the leading cause of acute abdominal pain, followed by cholecystitis and others.¹ The major complaint of surgical patients is the post-operative pain that occurs due to trauma to the tissues during surgery and activates pain pathways by the release of chemical mediators. Almost 22-67% patients experience moderate to severe pain during the first 24 hours after abdominal surgery as reported by various studies. The management of post-operative pain is the foremost challenge for surgeons as it is necessary for early mobilization and recovery as well as the psychological satisfaction of the patient.² Pain activates the sympathetic nervous system and renders the patient immobile. It also cripples the patient psychologically. All of this leads to more burden on the economy, hospitals, and healthcare

resources. In the modern era, post-operative pain relief is directed towards maximum comfort of the patient using safe and cost-effective analgesics.³

Diclofenac sodium belongs to the class of non-steroidal anti-inflammatory drugs (NSAIDs). They inhibit cyclooxygenase enzymes and prostaglandin synthesis. Diclofenac in particular inhibits both cyclooxygenase 1 and 2 enzymes as well as lipoxygenase and phospholipase A2. It is a potent analgesic for musculoskeletal and spasmodic pain.⁴ It has antipyretic effect and also helps in the reduction of inflammation.⁵

Diclofenac can be given orally as a tablet, through intramuscular injection or as a rectal suppository. Administration of the drug via the rectal route is easier and does not require a trained personnel as compared to the intramuscular route.⁶ Intramuscular administration should be done by following proper protocol & aseptic techniques, and failure to do so may result in localized infection or abscess formation.⁷ Administration of drug via the rectal route has the benefits of decreased first-pass metabolism, no gastric irritation, and higher local and systemic levels.⁸

Assessing the difference in pain relief and the need for rescue analgesia between diclofenac suppository and injectable diclofenac can guide clinicians in selecting the most effective and patient-friendly route of administration. This study aimed to compare the analgesic efficacy of diclofenac

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Muhammad Abu Usama
Postgraduate Resident, Department of Surgery
Mayo Hospital, Lahore
E-mail: ususama59@gmail.com

Received: October 3, 2025; Accepted: October 24, 2025

suppository versus injectable diclofenac in patients undergoing abdominal surgeries, to determine which route provides optimal post-operative pain control with reduced dependency on additional analgesics.

METHODOLOGY

This quasi-experimental study was conducted at the West Surgical Ward, Mayo Hospital, Lahore over a period of 8 months from June 2022 to January 2023 after taking approval from the ethical committee (Letter No. 233/RC/KEMU, 15-03-2021). A sample size of 136 participants was determined based on 80% power, 95% confidence interval, and the post-operative mean visual analog scale (VAS) pain scores for the diclofenac suppository and intramuscular injection groups reported as 3.88 ± 1.90 and 4.76 ± 1.75 , respectively.⁹ Patients were enrolled by using non-probability convenience sampling technique and were randomly divided into 2 groups of 68 each. Males and females, ≥ 18 years of age undergoing elective or emergency abdominal surgeries were included in the study. Patients with history of any previous abdominal surgery, NSAIDs contraindication (patients with gastritis, gastric or duodenal perforation), and patients on central nervous system (CNS) depressants or warfarin treatment were excluded.

Informed written consent was taken from all patients. The demographic data and body mass index (BMI) were recorded. Patients with BMI less 25 kg/m^2 were labeled as normal and those with $\geq 25 \text{ kg/m}^2$ were categorized as having deranged BMI. Group A was given diclofenac suppositories [Voltral suppository by DVAGO-100 mg twice daily (BD)] and group B was given intramuscular diclofenac [Injection Voren by DVAGO-75 mg or 1.5 mg/kg three times a day (TDS)]. If the pain was not relieved with diclofenac by either route, rescue analgesia was given in the form of injection nalbuphine via intravenous route at the dose of 10 mg.

Visual analog scale was used to measure pain at 0, 8, 16, and 24 hours post-operatively. Patients graded the severity of pain on VAS from 0 (no pain) to 10 (severe pain).⁹ The proportion of patients undergoing any adverse event (gastrointestinal tract discomfort, headache, itching) and requiring rescue analgesia were also recorded.

STATISTICAL ANALYSIS

Data entry and analysis was done by using Statistical Package for the Social Sciences (SPSS) version 26. Quantitative variables such as age, body mass index

(BMI), and VAS were presented as mean $\pm$ standard deviation (SD). Qualitative variables such as gender and adverse events were presented by frequency & percentages. The results for quantitative and qualitative variables were compared by using independent sample t-test and chi-square, respectively. A p-value < 0.05 was taken as significant.

RESULTS

The mean age of patients in group A and group B was 33.22 ± 9.58 years and 34.05 ± 10.70 years, respectively. The mean BMI of patients in group A was $25.71 \pm 2.35 \text{ kg/m}^2$ and the mean BMI of group B patients was $26.38 \pm 2.70 \text{ kg/m}^2$. Both groups didn't significantly differ in terms of age, gender, and BMI distribution ($p > 0.05$). In group A, only 15(11.1%) patients needed rescue analgesia as compared to 39(28.7%) patients in group B ($p = 0.001$). Also, 35(25.7%) patients faced adverse events in group B as compared to group A, where only 11(8.1%) patients suffered from adverse events ($p = 0.001$) (Table 1).

The patients receiving diclofenac suppositories showed significantly better pain relief compared to those who received Injection diclofenac via I/M route. Group A showed a reduction in mean pain scores at 0, 8, 16, and 24 hours post-operatively as compared to group B ($p < 0.001$) (Table 2). Pain reduction was greater in group A across all age categories. Participants aged 18-35 years had lower mean pain scores in group A (3.86 ± 0.89) compared to group B (4.13 ± 1.10) ($p = 0.028$). Similarly, in those aged ≥ 35 years, group A (3.75 ± 1.09) reported significantly less pain than group B (4.17 ± 1.27) ($p = 0.0043$). Both males and females in group A had significantly lower mean pain scores compared to their counterparts in group B ($p < 0.05$). The analgesic effect of group A treatment remained superior regardless of side effects. Among participants who experienced adverse effects, group A (3.75 ± 1.08) reported lower pain than group B (4.38 ± 1.16) ($p < 0.001$). The same pattern was observed in those without adverse effects ($p < 0.001$). Participants who required rescue analgesia had higher pain scores, but group A (3.85 ± 1.12) still showed significantly lower pain than group B (4.65 ± 1.30) ($p < 0.001$). Even among those not needing rescue analgesia, group A had better pain control ($p < 0.001$). When stratification was done for BMI, patients with normal & deranged BMI in group A had significantly lower mean pain score as compared to group B (p-value < 0.001) (Table 3).

Table 1: Demographic and Clinical Parameters of Groups A & B

Variables		Group A Diclofenac Suppositories (Frequency & Percentage)	Group B Injection Diclofenac (Frequency & Percentage)	Total	p-value
Age Groups (Years)	18 to <35	25(18.4%)	30(22.1%)	55(40.5%)	0.38
	≥35	43(31.6%)	38(27.9%)	81(59.5%)	
	Total	68(50%)	68(50%)	136(100%)	
Gender	Male	46(33.8%)	49(36.1%)	95(69.9%)	0.57
	Female	22(16.2%)	19(13.9%)	41(30.1%)	
	Total	68(50%)	68(50%)	136(100%)	
Adverse Events	Yes	11(8.1%)	35(25.7%)	46(33.8%)	0.001*
	No	57(41.9%)	33(24.3%)	90(66.2%)	
	Total	68(50%)	68(50%)	136(100%)	
Patients who Needed Rescue Analgesia	Yes	15(11.1%)	39(28.7%)	54(39.7%)	0.001*
	No	53(38.9%)	29(21.3%)	82(60.3%)	
	Total	68(50%)	68(50%)	136(100%)	

*Significant p-value

Table 2: Comparison of Post-Operative Mean Pain Scores between Study Groups

Time (Hours)	Group A (Mean Pain Scores)	Group B (Mean Pain Scores)	p value
0	4.94±0.96	5.67±0.79	<0.001*
8	4.4±0.99	5.25±1.09	<0.001*
16	3.95±0.89	5.06±1.53	<0.001*
24	3.75±1.09	4.69±1.05	<0.001*

*Significant p-value

Table 3: Comparison of Mean Post-Operative Pain Scores between Groups A & B regarding Demographic & Clinical Variables

Variables		Group A (Mean Pain Scores after 24 Hours (Mean±SD))	Group B (Mean Pain Scores after 24 Hours (Mean±SD))	p-value
Age Groups (Years)	18 to <35	3.86±0.89	4.13±1.1	0.028*
	≥35	3.75±1.09	4.17±1.27	0.004*
Gender	Male	3.7±1.07	4.16±1.07	0.0004*
	Female	3.88±1.16	4.46±1.08	<0.001*
Adverse Effects	Yes	3.75±1.08	4.38±1.16	<0.001*
	No	3.6±1.0	4.55±1.28	<0.001*
Rescue Analgesia	Yes	3.85±1.12	4.65±1.3	<0.001*
	No	3.58±1.1	4.49±1.32	<0.001*
BMI (kg/m ²)	Normal (18.5-24.9)	3.65±1.43	4.49±1.35	<0.001*
	Deranged (≥25)	3.86±0.51	4.62±1.1	<0.001*

*Significant p-value

DISCUSSION

The foremost challenge after a surgery is effective relief in pain with minimum possible side effects and cost-effectiveness.¹⁰ This study aimed to compare the effect of diclofenac suppositories and injection in the management and relief of pain after abdominal surgeries.

The mean age of patients in group A and group B was 33.22±9.58 years and 34.05±10.70 years, respectively. Pain reduction was significantly greater in group A in both age groups as compared to group B. Aleem et al. also reported that the mean age of patients who received diclofenac suppositories was 43.50±5.72 years and those who received diclofenac injections was 44.16±5.31 years. They reported

better pain control and relief with diclofenac suppositories; the most significant differences were recorded at 24 hours ($p=0.001$).¹¹ However, in our study, the most significant difference in score has been recorded at 16 hours, followed by 24 hours; this can be attributed to the difference in mean age of the populations in these studies.

In our study, 69.9% of the participants were males, whereas 30.1% were females. Pain reduction was significantly greater in group A across both gender categories as compared to group B. A previous study conducted in Faisalabad included approximately two-third males and one-third females. The study showed that diclofenac suppository was more effective at 12th hour ($p < 0.001$) as compared to 2nd and 6th hour in relief of post-operative pain.¹² A study conducted on pediatric patients with mean age of 6.05 ± 2.33 years for the relief of post-operative pain showed that patients who received diclofenac suppositories had lower pain scores ($p\text{-value}=0.03$) on VAS which corresponded to our study results.¹³ Another study was conducted to assess pain relief in the first 24 hours after cesarean section delivery. The study results showed better pain relief with diclofenac suppositories, similar to our study despite the fact that the sample population consisted of females only.¹⁴ Rab et al. included 72.45% males and showed effective pain relief with diclofenac suppositories after abdominal surgery.¹⁵ Another study compared the effects of the two forms of the drug in relief of the pain after vaginal delivery. The study showed that the patients who received diclofenac suppositories reported mild to moderate pain, while a few patients who received injection diclofenac reported severe pain as well, besides others who had mild to moderate pain ($p\text{-value}=0.002$ at 6 hours).¹⁶

In our study, the mean BMI of patients in group A & B was not significantly different. However, when stratification was done for BMI, patients with normal & deranged BMI in group A had significantly lower mean pain score as compared to group B ($p\text{-value} < 0.001$). A study conducted in Iran compared the effect of diclofenac suppositories with placebo group in relief of post-operative pain in patients undergoing laparoscopic cholecystectomy. Similar to our study, the placebo and diclofenac group did not significantly differ in terms of BMI distribution. The study showed significant reduction in mean pain scores on VAS of the patients who received diclofenac suppositories as compared to the other group ($p\text{-value} < 0.05$) but they did not reported the association of post-operative pain scores and BMI categories.¹⁷ Saeed et al. conducted a study on

females undergoing gynecological and obstetrical surgeries and reported that rectal diclofenac suppositories were effective for post-operative pain relief in patients with any BMI category.³

In our study, the results showed that mean pain on VAS was significantly lower for those who received diclofenac suppositories as compared to those who were given injection diclofenac at 0, 8, 16, and 24 hours ($p\text{-value} < 0.0001$). Altaf et al. compared the effectiveness of diclofenac suppositories versus injections in pain relief after perineal repair in primipara. Mean pain score on VAS in the suppositories group was 2.08 ± 0.96 and the score in the injection group was 5.92 ± 1.08 ($p\text{-value} < 0.0001$).¹⁸ A study compared rectal diclofenac suppository and transdermal diclofenac patch in the relief of pain after open cholecystectomy. The study concluded that the suppository form of the drug is more effective than transdermal patch ($p\text{-value} < 0.05$).¹⁹ These findings strongly support the results of our study. A study conducted in Nigeria compared rectal and intramuscular diclofenac for pain relief after abdominal surgery. The results showed significantly lower pain scores on VAS at 24 hours in the rectal suppository group. A statistically significant difference was found between the two groups ($p\text{-value} < 0.001$).²⁰ Elahabadi et al. conducted a study in Iran in which they compared diclofenac with acetaminophen suppositories in post-hemorrhoidectomy pain relief. Diclofenac suppositories had better analgesic effect and showed more satisfactory results.²¹ A study comparing oral and rectal diclofenac in pain relief after episiotomy and concluded that suppositories are far better in pain control. The mean pain score in the suppositories group was 2.93 as compared to 3.98 in the oral diclofenac group.²²

In our study, significant results showed that fewer (11.1%) patients in group A needed rescue analgesia as compared to 28.7% patients in group B. Hosny et al. conducted a study for relief of post-episiotomy pain and reported that only 1.1% patients ($p\text{-value}=0.01$) required rescue analgesia.²³ This was proportionately less as compared to our study and may be due to less extensive procedure and hence requiring lesser need for analgesia.

CONCLUSION

Rectal suppositories of diclofenac were significantly more effective for post-operative pain relief among patients undergoing abdominal surgeries as compared to diclofenac intramuscular injection. Diclofenac suppositories had shown significantly better pain relief as measured by VAS mean scores

at 0, 8, 16, and 24 hours. It was also observed that a smaller number of patients in the diclofenac suppositories group required rescue analgesia while more than half patients in the injection group demanded for rescue analgesia.

LIMITATIONS & RECOMMENDATIONS

The few limitations of this study included a single study setting and non-probability convenience sampling technique. Pain measurement by VAS introduced the risk of measurement bias since it is subjective and may vary from person to person. Multi-centered RCT with blinding should be done to further ascertain the results of this study. The groups should be further subdivided and matched for severity and type of abdominal surgery.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

M.A.U: Conceived the study, drafted the manuscript & coordinated data collection

F.L: Proofread the manuscript & ensured reference accuracy

F.M: Supervised the research & reviewed the manuscript

M.F: Performed statistical analysis & contributed to results interpretation

S.S.Z.J: Managed patient enrollment & clinical data verification

H.Z.A: Assisted in literature review & data organization

REFERENCES

- Sharma R, Lonare SB, Singh S, Al-Dwla H, Ranjan R. Clinical presentations and surgical outcomes in patients with nontraumatic acute abdominal pain: a prospective analysis. *Cureus*. 2024; 16(12):e75772. doi:10.7759/cureus.75772.
- Mehari EE, Bizuneh YB, Fentie DY, Arefayne NR. Prevalence and factors associated with acute postoperative pain after emergency abdominal surgery. *TOPAINJ*. 2022; 15. doi:10.2174/18763863-v15-e2208250.
- Saeed R, Mahboob A, Ali SM, Latif A, Ammar T, Kakepotto IA. Effectiveness of postoperative analgesia with rectal placement of diclofenac sodium in patients undergoing gynecological and obstetrical surgeries under general anesthesia. *Life and Science*. 2024; 5(1):120-5. doi:10.37185/LnS.1.1.515.
- Ede EE, Olaleye AA, Irechukwu JC, Nwaedu UN, Ejikeme BN, Ali VC, et al. Adjuvant rectal diclofenac for post operative analgesia after caesarean section - a randomized controlled study. *Open J Obstet Gynecol*. 2024; 14(4):591-609. doi:10.4236/ojog.2024.144052.
- Giri A, Palikhey A, Sah AK, Upadhyay HP, Giri BL. Analgesic efficacy of diclofenac and ibuprofen/paracetamol combination in post extraction cases. *J Coll Med Sci Nepal*. 2023; 19(1):32-7. doi:10.3126/jcmsn.v19i1.52792.
- Irechukwu JC, Eleje GU, Iwe BC, Ikpeze OC, Ikeotuonye AC, Ede EE, et al. A randomized controlled trial of rectal versus intramuscular diclofenac for post-operative analgesia after open myomectomy. *Eur J Obstet Gynecol Reprod Biol*. 2024; 292:244-50. doi:10.1016/j.ejogrb.2023.11.029.
- Altaf B, Kashif S, Sial SS, Noreen H, Pervaiz E, Awan SM. Comparison of intramuscular versus rectal diclofenac sodium in post caesarean pain relief. *Pak Armed Forces Med J*. 2022; 72(6):1982. doi:10.51253/pafmj.v72i6.7153.
- Okam PC, Ikechebelu JI, Eleje GU, Albert IC, Okpala BC, Okam CF, et al. Efficacy and safety of diclofenac suppository for postoperative pain relief after diagnostic hystero-laparoscopy and dye test: a double-blind, placebo-controlled, randomized trial. *Eur J Obstet Gynecol Reprod Biol*. 2024; 23:100326. doi:10.1016/j.eurox.2024.100326.
- Zulfiqar M, Ashraf M, Khan MIH, Naeem F, Shakeel S. To compare efficacy of diclofenac by intramuscular route and rectal route in post surgical pain. *Professional Med J*. 2021; 28(09):1269-75. doi:10.29309/TPMJ/2021.28.09.5885.
- Dieu A, Huynen P, Lavand'homme P, Beloeil H, Freys SM, Pogatzki-Zahn EM, et al. Pain management after open liver resection: procedure-specific postoperative pain management (PROSPECT) recommendations. *Reg Anesth Pain Med*. 2021; 46(5):433-45. doi:10.1136/rapm-2020-101933.
- Aleem S, Sarwar N, Naseem S, Abbas MA, Hameed S, Aleem A. Effectiveness of intramuscular diclofenac sodium injection versus diclofenac suppositories in post-operative pain management in inguinal hernia. *Pak J Med Health Sci*. 2022; 16(03):197-8. doi:10.53350/pjmhs22163197.
- Habib A, Iqbal MS, Choudry ZA, Lodhi MFB, Siddique R, Haider M. Comparison of post-operative pain with diclofenac rectal suppository vs intravenous tramadol in patients undergoing inguinal hernioplasty. *APMC*. 2024; 18(1):64-7. doi:10.29054/apmc/2024.1554.
- Bashir Z, Islam MN, Rahman MA, Islam SA, Sarker PC. Postoperative pain relief in pediatric surgery patients: effect of intravenous paracetamol in comparison with diclofenac suppository. *J Bangladesh Soc Anaesthesiologists*. 2022; 35(2):37-42. doi:10.3329/jbsa.v35i2.67894.
- Khan S, Majeed N, Ruqya, Mehdi M, Safdar F, Bibj S. Comparison of analgesic efficacy of diclofenac suppository with intramuscular diclofenac sodium in post-operative pain relief after cesarean delivery in the first 24 hours. *JRMC*. 2022; 26(4). doi:10.37939/jrhc.v26i4.1994.
- Rab A, Parveen S, Ali A, Nawaz A, Gul S, Musadaq M. Comparison of diclofenac suppository with injection tramadol in postoperative pain: diclofenac suppository with injection tramadol in postoperative pain. *PBMJ*. 2022; 5(6):46-9. doi:10.54393/pbmj.v5i6.591.
- Nafees R, Safdar Z, Baig FS, Zaib S, Iftikhar A. Rectal diclofenac: an effective modality for pain relief after vaginal birth. *Biomedica*. 2022; 38(1):39-43. doi:10.51441/BioMedica/5-604.

17. Asheghvatan A, Ahmadi Z, Kakaei F, Khodayari M, Ziaee M, Arjmand A. Pre-emptive effects of preoperative diclofenac suppository on pain management in patients undergoing laparoscopic cholecystectomy: a case-control study. *APTJ*. 2023; 3(1):1-8. doi:10.18502/aptj.v3i1.12486.
18. Altaf U, Asad AZ, Ashraf S, Tayyab M. Oral versus rectal diclofenac sodium efficacy in pain control after perineal repair in a primipara. *Esculapio - JSIMS*. 2023; 14(1):19-22. doi:10.51273/esc18.71415.
19. Agrawal P, Singh S, Gautam V, Khan BH. Efficacy of diclofenac transdermal patch versus diclofenac rectal suppository for management of postoperative pain following open cholecystectomy: a randomised clinical trial. *J Clin Diagn Res*. 2022; 16(2):FC01-4. doi:10.7860/JCDR/2022/53462.15969.
20. Uche-Omovoh IC, Ikeotuonye A, Nweke AN, Ogbo UE, Marcel-Onwudiwe CS, Oru CO, et al. Rectal versus intramuscular diclofenac for pain relief following elective caesarean section at Alex Ekwueme Federal University Teaching Hospital Abakaliki: a randomized controlled trial. *Asian J Med Health Sci*. 2025; 25(2):22-9. doi:10.5897/AJMHS2025.0283.
21. Elahabadi I, Khazaei A, Kamiab Z, Taghizadeh M, Bazmandegan G. Comparison of the effects of diclofenac and acetaminophen suppositories on post hemorrhoidectomy pain. *SJKUMS*. 2021; 26(4):50-8. doi:10.52547/sjku.26.4.50.
22. Mushtaq I, Kalsoom S, Nisa K, Jamil M, Safdar F, Majeed N. Oral versus rectal diclofenac sodium: comparison of efficacy for perineal pain relief after episiotomy. *Isra Med J*. 2022; 14(3):104-7. doi:10.55282/imj.oa1329.
23. Hosny M, ElNashaar D, Abdel Mageed O. Rectal versus oral diclofenac sodium in relieving post-episiotomy pain: randomized controlled study. *EBWHJ*. 2023; 13(3):242-8. doi:10.21608/ebwhj.2023.208634.1246.



Association of Radial Artery Diameter with Radial Artery Spasm in Patients Undergoing Transradial Cardiac Catheterization

Ismail Ahmed Khan, Umer Shafiq, Sohail Aziz, Zohair Aziz, Wajahat Lodhi, Waleed Hassan, Asif Manzoor

ABSTRACT

Objective: To assess the association of radial artery spasm (RAS) with radial artery diameter and other factors in patients undergoing transradial cardiac catheterization.

Methodology: This descriptive, cross-sectional study was conducted at the Fauji Foundation Hospital, Rawalpindi from June to September 2025 after ethical approval. After taking informed written consent, 150 patients who underwent coronary angiography or percutaneous coronary intervention using transradial approach were included. Prior to the procedure, radial artery diameter was assessed by the linear probe of Xario 100g Ultrasound machine. Radial artery spasm was assessed clinically by the operator and radial artery angiograms. Patients were divided into two groups: group I with no RAS and group II with RAS. The association of RAS with the radial artery diameter and other variables were determined. Statistical Package for the Social Sciences (SPSS) version 26 was used for data analysis.

Results: Radial artery spasm occurred in 35(23.3%) patients. The time to vascular access was significantly higher (150.60 ± 49.77 seconds) in patients with RAS as compared to those without spasm (43.69 ± 29.61 seconds) ($p=0.001$). The number of procedure attempts, episodes of vasovagal syncope (3.3% versus 0%) and access site crossover (6% versus 0%) were also significantly higher in patients with RAS. The radial artery spasm had a significant positive association with radial artery diameter, ejection fraction, angiographic findings, and treatment plan. The ROC curve showed an area under the curve of 0.876, showing the excellent ability of radial artery diameter to predict RAS.

Conclusion: Radial artery spasm affects a significant proportion of patients (23.3%) undergoing transradial cardiac catheterization. The radial artery diameter, vascular access time, need for ultrasound assistance to secure vascular assistance, additional vasodilator & glide wire, vasovagal syncope, and access site crossover were significantly higher in patients with RAS.

Keywords: Radial Artery. Cardiac catheterization. Percutaneous coronary intervention. Coronary angiography.

INTRODUCTION

Cardiac catheterization is an important and potentially lifesaving diagnostic and therapeutic modality, with the purpose to diagnose and treat conditions like coronary artery disease (CAD). It includes procedures such as diagnostic coronary angiography and percutaneous coronary intervention (PCI).¹ Various access sites can be used for left heart catheterization, such as the radial artery, ulnar artery, and femoral artery. Currently, the preferred site used to access in diagnostic and therapeutic left heart catheterization procedures is radial artery. This approach is also utilized in diagnostic coronary angiography and PCI. This is because of ease of access, patient comfort, reduced risk of blood vessels related complications, shorter hospital stay, and early patient mobilization proven by various clinical trials.² However, intraprocedural and postprocedural complications can still occur via transradial access. Intraprocedural

complications are radial artery spasm, kinking of the catheter, arterial dissection & perforation and vasovagal syncope. On the other hand, postprocedural complications include occlusion of the radial artery, pseudoaneurysm, hematoma formation, arteriovenous fistula, and nerve damage. These complications can potentially contribute to morbidity and mortality. They can lead to patient discomfort, limb dysfunction, long procedure duration, high chances of access site crossover, and life-threatening complications such as compartment syndrome.³

Radial artery spasm is generally identified when the operator feels resistance while advancing the sheath or catheter through this artery, causing pain in the forearm or arm. This complication is very common and is one of the leading causes of change in access site. Radial artery spasm occurs frequently owing to the vasoreactivity of the vessel. Most of the cases are mild that can be prevented and treated. Pain relief and sedation can markedly decrease the incidence of RAS. Proper selection of equipment also reduces the chances of RAS.⁴

The risk factors of RAS include females, young or older age, patients who are diabetic, hypertensive, dyslipidemic, smokers, and those with anxiety, and tachycardia are at risk of developing RAS. Radial artery spasm has been associated with both features

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Ismail Ahmed Khan
Fellow Interventional Cardiology, Department of Cardiology
Fauji Foundation Hospital, Rawalpindi
E-mail: xperiasp17@gmail.com

Received: September 29, 2025; Accepted: October 15, 2025

of radial artery and procedural factors. Small radial artery caliber and a weak radial pulse are radial artery parameters associated with RAS. Emergency procedures, multiple attempts, use of multiple catheters, and large sheaths can lead to RAS.⁵

The association of RAS with clinical parameters such as co-morbidities have been studied in detail, there is paucity of local data regarding the association of radial artery diameter with radial artery spasm, and while logic dictates that smaller radial artery diameter should be a significant contributor to this rather common complication. This study aimed to bridge this gap in knowledge that can potentially serve to identify patients at risk of radial artery spasm and its subsequent vascular complications, and hence reduce morbidity and mortality in patients undergoing transradial cardiac catheterization.

METHODOLOGY

The current descriptive, cross-sectional study was carried out at the Fauji Foundation Hospital, Rawalpindi from June to September 2025. After ethical approval (Letter No. 977/RC/FFH/RWP, 23-06-2025), the sample size of 150 was estimated using 11% frequency of radial artery spasm, 95% confidence interval, and 5% margin of error.⁶ The patients were included using non-probability convenience sampling.

Patients who underwent coronary angiography or PCI using transradial approach were included in the study. Exclusion criteria were those patients with absent radial pulse, prior abnormal modified Allen's test, history of radial artery occlusion, known radial artery anomalies, arteriovenous fistulas involving the radial artery, peripheral artery disease involving upper limbs, history of vascular complications in prior transradial cardiac catheterization, hemodynamic instability, and patients undergoing procedure from an alternative access site. Informed written consent was taken from the patients and their demographics, co-morbidities, electrocardiographic changes, and ejection fraction were noted.

The patency of the radial artery and the adequacy of collateral supply from the ulnar artery were checked through Modified Allen's test. After occluding both arteries and then releasing the ulnar artery, return of hand color within ≤ 5 seconds indicated adequate collateral flow and a normal result.⁷ Patients with abnormal test results were excluded from the study as per the defined exclusion criteria.

Prior to the procedure, radial artery diameters were assessed by the linear probe of Xario 100g Ultrasound machine, set on vascular mode at a frequency of 10 MHz. Measurements were taken in

cross-sectional plane in 2D mode at optimal gain settings, at a depth of 2-3 centimeters, keeping the arterial lumen in the middle third of the screen, with the calipers positioned in a manner to measure the anterior-posterior diameter from inner wall to inner wall. All measurements were taken 1cm proximal to the styloid process. As per the hospital's cardiac catheterization lab protocols, 5000 IU of I/V unfractionated heparin along with 200 mcg of I/V nitrates mixed in 10 ml of normal saline was injected via the sheath after securing the vascular access.

Radial artery spasm was clinically assessed by the operator performing the procedure based on the presence of pain or resistance during the manipulation of the sheath, catheters, or guidewires. In such instances, radial artery angiograms were performed to document radial artery spasm and differentiate it from other possible causes of pain and resistance, such as radial artery loops or tortuosity.⁵ The total number of attempts, time, and need of ultrasonographic (USG) assistance in securing vascular access, fluoroscopy time and coronary angiographic findings were recorded. Similarly, in the event of radial artery spasm, the need of additional vasodilators, the use of hydrophilic guidewires, or the need and location of access site crossover were also recorded. Any procedural complication, such as vasovagal syncope was also noted.

To eliminate interobserver bias, modified Allen's test, assessment of radial artery diameter on ultrasound and assessment of radial artery spasm were done by the consensus of 2 interventional cardiologists with experience of more than 200 transradial cardiac catheterizations per year.

STATISTICAL ANALYSIS

Data was analyzed using the Statistical Package for the Social Sciences (SPSS) version 26. Categorical and numerical variables were displayed as frequency (percentage) and mean (standard deviation), respectively. Independent t-test was used for the association of numerical variables. Chi-square and Fisher's exact test was used to analyze categorical variables. The receiver operating characteristic (ROC) curve was drawn for radial artery diameter to evaluate its ability to predict radial artery spasm. The area under the curve (AUC) of greater than 0.6 is considered meaningful, graded as poor if >0.6 but <0.7 , fair if >0.7 but <0.8 , good if >0.8 but <0.9 , and excellent if >0.9 .⁸ The cut-off value of radial artery diameter was estimated using the Youden Index. The Youden index (J) is used to

calculate the threshold value of a parameter with maximum sensitivity and specificity using the formula: sensitivity + specificity – 1.⁸ The results were considered significant at the p-value of <0.05.

RESULTS

Mean age of the patients was 59.21 ± 10.52 years with a range of 26 to 89 years. There were 115(76.7%) females and 35(23.3%) males. Most of the patients (42.6%) had stable angina, 26.7% had non-ST-elevation myocardial infarction (NSTEMI), 24% had ST-elevation myocardial infarction (STEMI), and 6.7% had unstable angina. Seventy seven (51.3%) patients were diabetic, 115(76.7%) were hypertensive, 9(6%) were smokers, and 22(14.7%) had a history of prior PCI. Most of the patients had triple vessel coronary artery disease (TVCAD) (26%) and double vessel coronary artery disease (DVCAD) (26%), followed by single vessel coronary artery disease (SVCAD) (18.7%). The majority of the patients were managed with medical treatment (38%), followed by PCI (36.7%).

Mean radial artery diameter of patients was 2.25 ± 0.35 mm. The mean angiographic fluoroscopic time was 3.05 ± 3.45 minutes. In patients who underwent PCI, the mean PCI fluoroscopic time was 9.89 ± 6.63 minutes, and in these patients, the total fluoroscopic time was 12 ± 6.7 minutes.

Radial artery spasm occurred in 35(23.3%) patients. The mean time to vascular access was significantly higher (150.60 ± 49.77 seconds) in patients with RAS as compared to those without spasm (43.69 ± 29.61 seconds). Whereas the angiographic, PCI, and total

fluoroscopic time were not statistically different ($p > 0.05$). A significantly higher proportion of patients with RAS required ultrasound assistance for vascular access (10.6%), use of additional vasodilators (18%), guidewire insertion (14.7%) and crossover to other site (6%) compared to those without RAS. The frequency of RAS also increased with increasing number of procedure attempts. Moreover, episodes of vasovagal syncope (3.3%) were also significantly higher in patients with RAS ($p = 0.001$) (Table 1).

The radial artery spasm had a significant positive association with its diameter. The mean radial artery diameter was much lower in patients with radial artery spasm. The ejection fraction was significantly lower in patients with spasm than with no spasm. Triple vessel disease was significantly associated with RAS (30.8%), followed by DVCAD (25.6%) and SVCAD (21.4%). The patients who underwent PCI had the highest proportion of RAS (29.1%) as compared to coronary artery bypass grafting (CABG) and medical treatment. The radial artery spasm had no significant link with age, gender, diabetes mellitus, hypertension, smoking, and prior PCI ($p > 0.05$). (Table 2).

The ROC curve drawn for radial artery diameter to predict radial artery spasm showed an area under the curve of 0.876. This shows that the radial artery diameter has an excellent ability to predict radial artery spasm. The cut-off value of radial artery diameter for predicting RAS was estimated to be 2.05 mm. At this cut-off value, the sensitivity of radial artery diameter was 77% and the specificity was 88% (Figure 1).




Figure 1: ROC Curve for Radial Artery Diameter to Predict Radial Artery Spasm

Table 1: Procedural Outcomes in Patients with or without Radial Artery Spasm

Parameters		Radial Artery Spasm (Frequency & Percentage)		Total	p-value
		No	Yes		
Number of Attempts	1	102(68%)	11(7.3%)	113(75.3%)	0.001*
	2	11(7.3%)	14(9.4%)	25(16.7%)	
	3	2(1.3%)	8(5.4%)	10(6.7%)	
	4	0(0%)	2(1.3%)	2(1.3%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Need for USG Assistance to Secure Vascular Access	No	111(74%)	19(12.7%)	130(86.7%)	0.001*
	Yes	4(2.7%)	16(10.6%)	20(13.3%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Need of Additional Vasodilator	No	115(76.7%)	8(5.3%)	123(82%)	0.001*
	Yes	0(0%)	27(18%)	27(18%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Need of Guide wire	No	112(74.7%)	13(8.6%)	125(83.3%)	0.001*
	Yes	3(2%)	22(14.7%)	25(16.7%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Vasovagal Syncope	No	115(76.7%)	30(20%)	145(96.7%)	0.001*
	Yes	0(0%)	5(3.3%)	5(3.3%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Crossover	No	115(76.7%)	26(17.3%)	141(94%)	0.001*
	Yes	0(0%)	9(6%)	9(6%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Time to Vascular Access (Seconds)	Mean±SD	43.69±29.61	150.60±49.77	-	0.001*

*Significant p-value

Table 2: Association of Radial Artery Spasm with Radial Artery Diameter and Clinical Parameters

Parameters		Radial Artery Spasm		Total	p-value
		No	Yes		
Radial Artery Diameter (mm)	Mean±SD	2.36±0.29	1.88±0.26	-	0.001*
Ejection Fraction	Mean±SD	50.96±9.93	45.86±10.87	-	0.01*
Angiographic Findings (Frequency & Percentage)	Normal Coronaries	19(12.7%)	4(2.6%)	23(15.3%)	0.001*
	Minor CAD	9(6%)	2(1.3%)	11(7.3%)	
	Subcritical CAD	9(6%)	1(0.7%)	10(6.7%)	
	SVCAD	22(14.7%)	6(4%)	28(18.7%)	
	DVCAD	29(19.3%)	10(6.7%)	39(26%)	
	TVCAD	27(18%)	12(8%)	39(26%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	
Treatment Plan (Frequency & Percentage)	Medical Rx	46(30.7%)	11(7.3%)	57(38%)	0.001*
	PCI	39(26%)	16(10.7%)	55(36.7%)	
	CABG	30(20%)	8(5.3%)	38(25.3%)	
	Total	115(76.7%)	35(23.3%)	150(100%)	

*Significant p-value

DISCUSSION

The frequency of RAS in our study was 23.3%. Rocznik et al. also reported symptomatic angiographic RAS in 24.3% of patients.⁹ This was higher than 11% reported in a previous study.⁶ The reported incidence of radial artery spasm ranges from 1% to 34%, while radial artery occlusion occurred in up to 19.7% of cases. This wide

variation may be explained by the differences in arterial size, operator experience, and the equipment used during the procedure.¹⁰ Patel et al. observed the RAS rate of 5.5% and emphasized that the type of vasodilator used in transradial access can also contribute to a wide difference in the occurrence of radial artery spasm.¹¹

The mean radial artery diameter of patients was 2.25 ± 0.35 mm in our study. Similarly, Hasan et al. observed that the mean radial artery diameter was 2.35 ± 0.41 mm. Moreover, literature showed that the diameter of the radial artery was smaller in Asians (2.63 ± 0.35 mm) as compared to the western population (3.6 ± 0.8 mm). The smaller caliber of the radial artery and its resultant tendency to develop spasms may lead to higher procedural failure in the transradial approach.¹²

In the current study, patients with RAS had a significantly smaller mean radial artery diameter (1.88 ± 0.26 mm vs. 2.36 ± 0.29 mm) ($p=0.001$). Sanketh et al. found that the frequency of radial artery occlusion was 24% in patients with small radial artery diameter (<2.5 mm) as compared to 6.7% in patients with larger diameter ($p=0.03$).¹³ Radial artery dimension was smaller among RAS patients as compared to those without RAS but the results were not significant in another study.⁹

The average patient age was 59.21 ± 10.52 years in our study and most of the patients were females (76.7%). The age and gender distribution had no significant association with RAS incidence. In another study, the average age was 62.5 ± 11.2 years with 62.3% males. The patients with smaller radial artery diameters were generally older and had a lower proportion of males. However, multivariate analysis revealed that age and gender were not significant predictors of radial artery occlusion.¹³ In a study by Mazhar et al., age was not significantly associated with RAS however, RAS showed a significant association with the female gender.⁶ Another study reported that the frequency of RAS was significantly higher among females (66.6%; $p=0.003$) and those aged ≤ 65 years (62.5%; $p=0.001$).¹⁴ Rocznik et al. also found that women were significantly more likely to experience radial artery spasm as compared to men (OR = 2.94, $p=0.02$).⁹

In our study, 76.7% of the patients were hypertensive and 51.3% were diabetic. Only 6% were smokers. The association of RAS with hypertension, diabetes mellitus, smoking, and prior PCI was insignificant. Previous studies didn't find any significant association of RAS with diabetes and hypertension.^{6,15} Hasan et al. also reported that radial artery diameter wasn't significantly linked to diabetes and hypertension.¹² In addition to diabetes

and hypertension, smoking was also not considered a predictor of RAS.¹⁴

Fluoroscopic time in our study did not differ significantly between patients with RAS and without RAS. Procedural factors like puncture attempts, longer vascular access time, more frequent use of ultrasound, guide wire, additional vasodilators, as well as higher rates of vasovagal episodes and access site crossover were strongly linked to RAS in our participants ($p \leq 0.001$). Another study reported that first-attempt failure significantly predicted spasm.⁹ Roy et al. reported increased procedural time and access site crossover in patients who developed radial artery spasm while undergoing transradial access.¹⁶ More than one procedural attempts were linked with RAS in a study conducted in 2023.¹⁵ Another study was conducted to compare radial artery diameter pre and post-radial coronary angiography and a change in radial artery diameter was significantly associated with the number of puncture attempts.¹⁷ Our data reinforce that minimizing puncture attempts can reduce spasm risk.

A majority of RAS patients needed additional vasodilators in our study. Vasodilators such as nitric oxide donors and calcium channel blockers can be administered individually or in combination with other agents to create an anti-spasmodic radial "cocktail" aimed at preventing or reducing radial artery spasm.¹⁰ Rocznik et al. used nitroglycerine and lidocaine cocktail but administration of the radial cocktail did not result in a statistically significant reduction in the odds of spasm ($p=0.48$).⁹

CONCLUSION

Radial artery spasm affects a significant proportion of patients (23.3%) undergoing transradial cardiac catheterization. The vascular access time, need for ultrasound assistance to secure vascular access, additional vasodilator & guide wire, vasovagal syncope, and access site crossover were significantly higher in patients with RAS. It had a significant association with small radial artery diameter, ejection fraction, angiographic findings, and treatment plan.

LIMITATIONS & RECOMMENDATIONS

The small sample size and conduction of the study at a single center affect the generalizability of the results. Future multicenter studies should be conducted on a large scale in Pakistan.

Small radial artery diameter is an associated factor of radial artery spasm. These patients must be

considered at high-risk and managed optimally to reduce the chances of RAS.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

I.A.K: Conceived and designed the study, supervised data collection, and critically reviewed the manuscript.

U.S: Contributed to study design, performed data analysis and interpretation, and drafted the manuscript.

S.A: Assisted in data collection, clinical evaluation, and literature review.

Z.A: Participated in patient selection, data acquisition, and manuscript formatting.

W.L: Contributed to data analysis and interpretation of results.

W.H: Assisted in statistical analysis and preparation of tables and figures.

A.M: Supervised the overall research process, reviewed the final manuscript for intellectual content, and approved it for publication.

REFERENCES

- Holzer RJ, Bergersen L, Thomson J, Aboulhosn J, Aggarwal V, Akagi T, et al. PICS/AEPC/APPCS/CSANZ/SCAI/SOLACI: expert consensus statement on cardiac catheterization for pediatric patients and adults with congenital heart disease. *JACC Cardiovasc Interv.* 2024; 17(2):115-216. doi:10.1016/j.jcin.2023.11.001.
- Kis M, Soydan E. Preservation of radial vasomotor functions through the anatomic snuffbox: a prospective comparison with other radial accesses during coronary angiography. *J Coll Physicians Surg Pak.* 2020; 30(11):1121-5. doi:10.29271/jcpsp.2020.11.1121.
- Safitri H, Alkatiri AA, Firman D, Mangkuanom AS. Managing transradial artery access complications during coronary intervention: three case reports. *JAPSC.* 2023; 2:e28. doi:10.15420/japsc.2022.29.
- Riangwiwat T, Blankenship JC. Vascular complications of transradial access for cardiac catheterization. *US Cardiol.* 2021; 15:e04. doi:10.15420/usc.2020.23.
- Zus AS, Crisan S, Luca S, Nisulescu D, Valcovici M, Patru O, et al. Radial artery spasm-a review on incidence, prevention and treatment. *Diagnostics (Basel).* 2024; 14(17):1897. doi:10.3390/diagnostics14171897.
- Mazhar MW, Tuyyab F, Samin A, Khan KA, Khan S, Ullah A, et al. Incidence and predictors of radial artery spasm during left heart catheterization. *Pak Armed Forces Med J.* 2020; 71(Suppl 2):S288-92. Available from: <https://www.pafmj.org/PAFMJ/article/view/7763>.
- Moon J, Lee GH, Park TJ, Ki SH. Is the modified Allen's test a useful tool for evaluating the vascular dominance of the forearm? *J Korean Med Sci.* 2025; 40(36):e230. doi:10.3346/jkms.2025.40.e230.
- Corbacioglu SK, Aksel G. Receiver operating characteristic curve analysis in diagnostic accuracy studies: a guide to interpreting the area under the curve value. *Turk J Emerg Med.* 2023; 23(4):195-8. doi:10.4103/tjem.tjem_182_23.
- Roczniak J, Tarnawski A, Dziewierz A, Glanowski S, Pawlik A, Sabatowski K, et al. Radial artery spasms - angiographic morphology, risk factors and management. *Postepy Kardiol Interwencyjne.* 2024; 20(1):53-61. doi:10.5114/aic.2024.136936.
- Coghill EM, Johnson T, Morris RE, Megson IL, Leslie SJ. Radial artery access site complications during cardiac procedures, clinical implications and potential solutions: the role of nitric oxide. *World J Cardiol.* 2020; 12(1):26-34. doi:10.4330/wjc.v12.i1.26.
- Patel K, Duckett M, Vidovich MI, Ibrahim K. Intra-arterial nicardipine versus verapamil during transradial access coronary catheterization. *Cardiovasc Revasc Med.* 2025; 74:40-4. doi:10.1016/j.carrev.2024.08.008.
- Hasan MM, Dutta AK, Al Mahmood MR, Kanungo S, Rahman A, Hassan MR. Quantitative angiographic radial artery diameter measurement and its relationship with common variables: A cross-sectional study. *Medicine (Baltimore).* 2025; 104(32):e43797. doi:10.1097/MD.00000000043797.
- Sanketh SJ, Charantimath NV, Kumar S. Relationship between radial artery diameters with incidence of radial artery occlusion after transradial coronary catheterization: an observational cohort study. *Sch J App Med Sci.* 2024; 12(11):1652-6. doi:10.36347/sjams.2024.v12i11.035.
- Khan M, Daud MY, Awan MS, Khan MI, Khan H, Yousuf MA. Frequency and predictors of radial artery spasm during coronary angiography/percutaneous coronary intervention. *J Ayub Med Coll Abbottabad.* 2020; 32(3):357-9. Available from: <https://jamc.ayubmed.edu.pk/index.php/jamc/article/view/7643/2936>.
- Curtis E, Fernandez R, Khoo J, Weaver J, Lee A, Halcomb L. Clinical predictors and management for radial artery spasm: an Australian cross-sectional study. *BMC Cardiovasc Disord.* 2023; 23(1):33. doi:10.1186/s12872-023-03119-9.
- Roy S, Kabach M, Patel DB, Guzman LA, Jovin IS. Radial artery access complications: prevention, diagnosis and management. *Cardiovasc Revasc Med.* 2022; 40:163-71. doi:10.1016/j.carrev.2021.12.007.
- Abdelrahman AO, Soliman MA, Elkersh AM, Mina MB. Angiographic evaluation of radial artery diameter pre-and post-coronary angiography. *EJHM.* 2023; 90(1):1365-71. doi:10.21608/ejhm.2023.282151.



Early Clinical Exposure through Modular Curriculum in Undergraduate Medical Students: Student Experiences

Lamia Yusuf, Aqsa Aslam, Maaz Ahmed, Muhammad Fayyaz Atif, Muhammad Hashim Khan, Muhammad Bin Ahsan

ABSTRACT

Objective: To explore experiences and perceptions of early clinical exposure (ECE) in undergraduate medical students under a modular curriculum through a mixed-methods approach.

Methodology: This cross-sectional mixed-methods study was conducted at Rashid Latif Khan University Medical College, Lahore from July to September 2025, following a recently introduced modular curriculum after ethical approval. The participants were second-year MBBS students who had completed at least one year of learning under the modular system with early clinical exposure components. After obtaining written, informed consent, 100 students were included using non-probability convenience sampling technique. Quantitative data was collected through a structured questionnaire using 5-point Likert scale assessing integration of basic and clinical sciences, motivation to learn, communication with patients, and overall satisfaction. Qualitative perceptions were obtained through focus group discussions (FGDs), transcribed verbatim and analyzed thematically using NVivo software. Quantitative data was analyzed using Statistical Package for the Social Sciences (SPSS) version 25.

Results: The majority of the students were of the opinion that ECE resulted in enhanced incorporation of basic and clinical sciences (mean Likert score 4.3/5), it also improved the motivation to learn (4.5/5), and better-quality communication skills with patients (4.2/5). Qualitative data analysis with thematic analysis discovered four major themes, i.e. (1) Linking theory and practice; students recognized the significance of learning in a specific context (2) Motivation through realistic exposure - clinical learning environment stimulated them for enhanced learning (3) Professionalism and communication - students felt an improvement in their self-confidence, communication skills and empathy with the patients; and (4) Challenges and barriers - despite major positive outcomes overcrowding, lack of time, and inconsistent faculty input posed difficulties.

Conclusion: The majority of the students perceived early clinical exposure within modular curricula positively. It enhanced students' motivation, understanding, and communication skills. Fixing the logistical and faculty training challenges can maximize the impact.

Keywords: *Clinical competence. Professionalism. Empathy. Curriculum.*

INTRODUCTION

Globally, medical education has shifted from traditional, discipline-based curricula to modular and competency-based frameworks for the undergraduate medical and dental students. Such an integrated curriculum incorporates basic science disciplines with clinical subjects from early years, allowing the bridging of knowledge with clinical application. The majority of such curriculum designs have a key feature of early clinical exposure (ECE), providing them a learning experience of patient care and communication from the initial years of medical education.¹

Early clinical exposure is believed and proven to bridge the gap between basic medical learning and actual patient interaction. It enhances students' motivation, understanding, and learning in clinical

contexts by providing them an exposure to clinical environments from the beginning. It not only enhances context specific learning and professionalism but also helps in early acquisition of communication skills and empathy among the medical students. Literature review suggests that a structured early clinical exposure improves self-assessed professional competence and confidence in clinical skills.² Studies conducted within Pakistan have almost similar findings, providing evidence of benefits to the undergraduate medical students perceiving it as a helpful tool of instruction in concept building and enhancing engagement during the pre-clinical years.³

Apart from bridging the gap between theory and practice, ECE has the advantage of contributing significantly to the holistic professional progress of medical students by building their professional identity and emotional endurance. Learning in clinical contexts indirectly inculcates empathy, ethical values, and reflective thinking. Such qualities are expected to be essential in a lifelong learner who is providing compassionate care.⁴ Early clinical exposure in a specific clinical environment not only helps the students in a better understanding of the complexities of healthcare systems but also provides

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Muhammad Fayyaz Atif
Professor, Department of Community Medicine
Rashid Latif Khan University Medical College, Lahore
E-mail: dr.fayyazatif@hotmail.com

Received: October 8, 2025; Accepted: October 30, 2025

insight to interprofessional collaboration, and social determinants of health in the local context. The opportunity of working and learning in an authentic clinical situation during early educational years provides an additional benefit of adopting the values and responsibilities of the medical profession by the students. Hence, it incorporates a sense of determination and belonging that provides continuous motivation throughout their training.⁵

The University of Health Sciences (UHS), Lahore has introduced the modular integrated curriculum 2k23 in 2023 in its affiliated medical colleges. Data pertaining to ECE is limited to the introduction of clinical lectures in the first two years, while the modular integrated curriculum 2k23 provides a real-time clinical interaction with students learning relevant clinical skills in a specific clinical setting.⁶ The responses across institutions are variable, and students' perception has not yet been explored in the local context. Therefore, we decided to conduct a mixed-method study to explore experiences and perceptions of the students undergoing ECE in the modular integrated curriculum 2k23. This mixed-method study seeks to explore the perceptions, experiences, and challenges of the undergraduate medical students undergoing ECE within the UHS modular integrated curriculum 2k23. By integrating quantitative survey data with qualitative insights, the study aims to provide a comprehensive understanding of how ECE contributes to learning, motivation, and professionalism in the early years of medical education. We believe that the conclusions from this research will not only provide guidelines to curriculum organizers but will also suggest measures for faculty development and capacity building. It will also assist the policymakers and degree awarding bodies in promoting the early clinical learning experiences across Pakistani medical schools.

METHODOLOGY

This cross-sectional mixed-methods study was conducted at Rashid Latif Khan University Medical College, Lahore from July to September 2025 that implemented a modular MBBS curriculum in 2023, after ethical approval (Letter No. RLKUMC/IRB/0063, 24-06-2025). The participants were second-year MBBS students who had completed at least one year of learning under the modular system with early clinical exposure components. After obtaining informed written consent, 100 students were included using non-probability convenience sampling technique.

Data was collected through a structured, pretested questionnaire. It was developed following AMEE Guide No. 87 for developing questionnaire, the literature review, development of relevant items, and preliminary questionnaire, pilot testing with revision and refining based upon the feedback of pilot testing. This was followed by pre-testing to validate the questionnaire. A Cronbach's alpha of 0.7 was obtained indicating internal consistency of the item. It was a structured questionnaire consisting of statements rated on a 5-point Likert scale (1=strongly disagree, 2=disagree, 3=neutral, 4=agree, 5=strongly agree). The survey covered four key domains of the ECE experience: integration of basic and clinical sciences, motivation to learn, communication with patients, and overall satisfaction.

In-depth qualitative insights were gathered through two FGDs with students (n=10 in each) enrolled through purposive sampling. Each FGD was facilitated using a semi-structured guide that encouraged students to discuss the benefits, challenges, and suggestions regarding early clinical exposure. Open-ended questions were designed to explore experiences corresponding to themes. These questions related to themes are given in Table 1.

Table 1: Qualitative Research Questions Aligned with Themes

Theme	Qualitative Research Question
Bridging Theory and Practice	In what ways has early clinical exposure helped you connect with what you learn in lectures with actual patient care?
Motivation through Real-Life Exposure	How did meeting real patients early in your training affect your motivation or interest in learning medicine?
Professionalism and Communication	Can you describe how early patient interactions have influenced your communication skills or professional behavior?
Challenges and Barriers	What difficulties or barriers have you encountered during your early clinical exposure sessions?

STATISTICAL ANALYSIS

Quantitative data was analyzed using the Statistical Package for the Social Sciences (SPSS) version 25. The descriptive data, including frequencies (percentages) and mean±standard deviation (SD) for each item and domain of the questionnaire (integration, motivation, communication, and overall satisfaction) were calculated. These are mentioned in Table 2. As the study population comprised only 2nd year MBBS students, the analysis focused on summarizing response trends rather than subgroup comparisons. To analyze the trends of students' perceptions, combined scores from each domain were calculated.

For qualitative data, NVivo software (version XX) was used. We followed the principles of Braun and Clarke for qualitative data analysis to generate themes and thematic analysis. Audio recordings from the two focus group discussions were transcribed verbatim and introduced into NVivo for coding and organization. Utilizing thematic analysis approach, repeated patterns and categories were identified inductively from the data (Figure 1). This was followed by iterative coding to draft emerging themes connected to students' perceived advantages, challenges, and proposals for improvement regarding early clinical exposure. For refining and validation of final themes, thematic maps and query

functions in NVivo were used to safeguard reliability and uniformity of interpretation.

RESULTS

Hundred students gave consistently positive ratings to their early clinical exposure. Most of them agreed that ECE improved their understanding of the relevance of basic sciences in a clinical context (82% agreement; mean Likert score 4.3/5), enhanced their motivation to learn (88% agreement; mean 4.5/5), and increased their confidence in communicating with patients (76% agreement; mean 4.2/5). In terms of overall impact, 80% of the students rated the ECE experience as beneficial to their education (Table 2).

Data examination through NVivo-assisted thematic analysis revealed that ECE not only improved integration between basic and clinical sciences, but it also improved student motivation and enhanced professional and communication skills. There were however, some challenges and contextual barriers that were identified by the students including overcrowding in clinical settings, time restraints, and unpredictable faculty guidance. In general, the findings in this study support the inclusion of structured and well-supervised ECE sessions in the early medical curriculum (Table 3).



Figure 1: Word Cloud Depicting Key Themes Emerging from Students Perspectives on Medical Education

Table 2: Mean Scores of Quantitative Domains of ECE

Domain	Description	Likert Score (out of 5) (Mean±SD)	Agree/Strongly Agree (Percentage)	Interpretation
Integration of Basic & Clinical Sciences	ECE enhanced the linkage between theory and practice	4.3±0.55	82%	Strongly positive
Motivation to Learn	ECE improved students' enthusiasm and engagement in learning	4.5±0.48	88%	Very positive
Communication Skills	ECE improved student–patient communication and confidence	4.2±0.6	76%	Positive
Overall Satisfaction with ECE Experience	ECE was perceived as a useful tool with positive impact on learning	4.33±0.54	80%	Positive

Table 3: Thematic Analysis of Qualitative Data

Theme	Subthemes (Identified Through NVivo Coding)	Representative Verbatim Comments
Linking Theory and Practice	Comprehending the significance of basic sciences through the incorporation of clinical examples	“When I saw the patient with anemia, I could eventually connect what we studied in physiology to the tangible problem. ECE facilitated me to understand why basic sciences actually matter.”
	Strengthened classroom learning leading to context-based learning	“I couldn’t understand the significance of studying physiology in so much detail in lectures unless I observed the clinical cases in ECE, everything started connecting in my mind”
	Augmented knowledge retention	“The anatomy of brachial plexus became so clear and easy when I saw a patient with a fractured humerus, I could visualize exactly which nerve was affected and why”.
Motivation Through Realistic Exposure	Enhanced internal motivation of students	“I felt like a real doctor in training. After the ECE sessions, I went home and read more because I wanted to know what I had seen.”
	Appealing, relevant, and meaningful learning	"When I learnt how to examine a patient's pulse and blood pressure, I instantly was eager to recall the physiology behind it. I wanted to comprehend what was happening inside the body."
	Clinical learning environment increased curiosity and self-study.	“After examining a patient with anemia in the ward, I felt inquisitive to understand the biochemical basis. For the first time, I read metabolism not for the test, but to connect it with real life.”
Professionalism and Communication	Enhanced self-confidence during patient interaction	"Initially, I was under confident to talk, but now I can communicate more confidently with patients. I realized that patients are people, not just cases, ECE taught me compassion."
	Development of empathy and professionalism	“When I saw patients with multiple health problems, I realized how much empathy matters & it made me think beyond books and marks.”
	Understanding the ethical aspects of care	“I learnt my first lesson about secrecy, patient confidentiality, and ethics during ECE, we were told not to discuss patient details outside the ward”.
Challenges and Barriers	Overcrowding in clinical settings; limited time for observation	"The clinical batches had too many students at one bedside resulting it difficult to learn. Small batches might help in a better understanding"
	Inconsistent faculty facilitation	“Each teacher had a different approach in clinical methodology, it got confusing sometimes.”
	Lack of structured debriefing	“There was generally no discussion on how we performed in ECE session. The sessions would have been more fruitful if there was a short debrief so we could clear our doubts and reflect on what we learned”

DISCUSSION

Early clinical exposure (ECE) is an educational approach that familiarizes medical students with patient interactions during the early stages of the MBBS program. It encourages students to enhance their academic knowledge, clinical abilities, and communication skills, thereby boosting their confidence.⁷

The findings of our study demonstrate that early clinical exposure serves as an effective pedagogical approach in bridging the gap between basic and clinical sciences in the early years of medical education. The higher mean Likert scores across all domains; motivation to learn, integration of basic & clinical sciences, communication skills, and overall satisfaction underscore the strong positive perception of students towards ECE. Similar outcomes have been reported in other studies. In a study by Oshiro et al., integrated modular curriculum in Tokyo led to improved self-assessment scores of medical students in various professional competencies. The students were able to correlate basic sciences with the practical knowledge, contributing to long-term retention of concepts and better efficacy.⁸ Another study from Ethiopia reported a positive role of ECE on the motivation of students to learn, problem-solving abilities, professional knowledge, and orientation towards the community.⁹ Ahire et al. stated that ECE improved students' understanding of concepts, enhanced knowledge retention, and increased interest in the subject.¹⁰ In a study, ECE positively influenced cognitive and affective learning domains. Students described a deeper comprehension of concepts of basic sciences when encountered in clinical contexts. The ability to visualize theoretical content in patient scenarios appeared to reinforce meaningful learning, aligning with the principles of constructivist and experiential learning theories. The sense of curiosity and intrinsic motivation described by students after ECE reflects self-directed learning behaviors, which is a key outcome desired in outcome-based medical curricula.¹¹ A study conducted at the Rawal Institute of Health Sciences, Islamabad assessed the effects of early clinical exposure in 3rd to final year MBBS students. The results of the study showed that ECE is associated with enhanced confidence, communication, and clinical skills of the students.¹² Similar to our study, a study revealed that 90% of the medical students were satisfied with the ECE program. Students reported that ECE facilitated adaptation to the clinical environment, increased confidence in

communicating with supervisors & patients, and enhanced their abilities to learn clinical skills.¹³

In our study, another important finding was the improvement in empathy and professional attitude among medical students. According to a study from Dubai, early patient exposure fosters emotional intelligence and nurtures empathy, allowing students to view patients as individuals rather than disease entities. It contributes significantly to the development of professionalism and ethical awareness in undergraduate medical students.¹⁴ In another study, ECE has also been shown to promote professional development in the initial stages of medical education.¹⁵ Ingale et al. reported that ECE helps the students to learn clinical skills by providing an engaging learning environment, cultivate empathy, foster a patient-focused approach to care, and develop professional attitudes such as the value of teamwork, effective communication, and ethical decision-making.¹⁶

However, our study identified that despite the overall positive impact, students found challenges such as overcrowding during sessions, inconsistent facilitation, and lack of structured debriefing. These issues are consistent with other studies. A study pointed out that ECE has complex dynamics depending on a large extent on student-teacher relationship. This is because the students consider their clinical teachers as mentors and role models. Insufficient guidance and poorly structured programs can hinder the effectiveness of ECE.¹⁶ Another study highlighted several barriers to implementing ECE, with increased workload of the clinical doctors as the most common challenge. Other significant obstacles included poor coordination between academic & healthcare institutions and inadequate orientation on the implementation process.⁹ A study reported a decline in student satisfaction with ECE, attributed to the absence of structured programs in hospitals.¹³

CONCLUSION

The majority of the students perceived ECE within modular curricula positively. Early clinical exposure, when implemented systematically with adequate faculty support, can enhance motivation to learn, integration of basic & clinical sciences, communication skills, and overall satisfaction in early years of medical students. It also contributes to improvement in empathy and professional attitude among medical students. Its incorporation into the preclinical curriculum of Pakistani medical colleges should be encouraged to promote active, reflective, and patient-centered learning.

LIMITATIONS & RECOMMENDATIONS

The study included only second-year MBBS students from a single medical institution. Future research should include students from different academic years and involve multiple medical colleges following UHS integrated curriculum to provide a more comprehensive evaluation of ECE-integrated curricula.

It is recommended that smaller group rotations, faculty development workshops, and structured debriefing sessions be implemented to address existing barriers and enhance the effectiveness of ECE in improving learning outcomes.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

L.Y: Concept and study design, data analysis, final proofreading, and approval of the manuscript.

A.A: Literature review, and questionnaire administration, data validation, results compilation, and critical manuscript review.

M.A: Data collection, statistical analysis and interpretation of quantitative data.

M.F.A: Data collection, facilitation of focus group discussions and qualitative data analysis.

M.H.K: Literature review, bibliography, data collection, and formatting.

M.B.A: Literature review, bibliography, data collection, and formatting.

REFERENCES

1. Datta A, Prabha AGT, Ananthakrishnan N. A comparative critical analysis of competency-based curricula prescribed by regulators for MBBS, BDS, and BSc Nursing programs in India. *Cureus*. 2025; 17(7):e89178. doi:10.7759/cureus.89178.
2. Kalusopa VM, Katowa-Mukwato P, Chitundu K, Mvula M, Nzala S, Kabinga-Makukula M, et al. Experiences of early and enhanced clinical exposure for postgraduate neonatal nursing students at the University of Zambia, School of Nursing Sciences: lessons and implications for the future. *Open J Nurs*. 2023; 13(6):352-67. doi:10.4236/ojn.2023.136024.
3. Ahmed A, Bibi A, Nisar N, Salman R, Yasmin R, Aslam U. Usefulness of clinical lectures in first two years of integrated teaching: students' perspective. *PJMHS*. 2022; 16(9):313-5. doi:10.53350/pjmhs22169313.
4. Tayade MC, Latti RG. Effectiveness of early clinical exposure in medical education: Settings and scientific theories - review. *J Educ Health Promot*. 2021; 10:117. doi:10.4103/jehp.jehp_988_20.
5. Krishnasamy N, Hasamnis AA, Patil SS. Developing professional identity among undergraduate medical students in a competency-based curriculum: Educators' perspective. *J Educ Health Promot*. 2022; 11:361. doi:10.4103/jehp.jehp_329_22.
6. Naseem N. The First year of successful execution & examination of integrated modular curriculum 2k23 by UHS Pakistan. *Pak J Med Sci*. 2024; 40(5):1047. doi:10.12669/pjms.40.5.9951.
7. Maheshwari KU. Role of early clinical exposure for clinical training among medical undergraduate students. *Int J Clin Biochem Res*. 2023; 10(3):184-6. doi:10.18231/ijicbr.2023.032.
8. Oshiro T, Suzuki S, Kagawa N, Ono H, Goto R, Furuta A, et al. Integration of early clinical exposure into curriculum enhances self-assessment of professional competencies in medical practice. *BMC Med Educ*. 2025; 25(1):1139. doi:10.1186/s12909-025-07678-7.
9. Ewnte B, Yigzaw T. Early clinical exposure in medical education: the experience from Debre Tabor University. *BMC Med Educ*. 2023; 23(1):252. doi:10.1186/s12909-023-04221-4.
10. Ahire PB, Kankhare SB, Nomulwar SG, Gaikwad A. Perception of students towards early clinical exposure in Anatomy. *EJCM*. 2025; 15(3):786-90. doi:10.5083/ejcm/25-03-136.
11. Singh RA. Perception of early clinical exposure (ECE) among phase I MBBS students in a medical college in Northeastern India. *JMSH*. 2024; 10(2):169-74. doi:10.46347/jmsh.v10.i2.24.132.
12. Aman K, Hassan B, Qamar Z, Riaz W, Gohar H, Abdullah S. Impact of structured early clinical exposure on confidence and competence among undergraduate health science students in Islamabad: clinical exposure on student confidence. *PJHS*. 2025; 6(6):149-53. doi:10.54393/pjhs.v6i6.3231.
13. Kachuei M, Rezazadeh M, Hosseinzadeh Davarzani M, Hashemi A, Mousavi AS. Satisfaction survey of early clinical exposure in medical students: a cross-sectional study. *Ann Med Surg*. 2025; 87(10):6295-302 doi:10.1097/MS9.0000000000003733.
14. Al-Jayyousi R, Abu Mahfouz N, Otaki F, Paulus A, Czabanowska K, Zaman Q, et al. Investigating the learning value of early clinical exposure among undergraduate medical students in Dubai: a convergent mixed methods study. *BMC Med Educ*. 2025; 25(638). doi:10.1186/s12909-025-07212-9.
15. Liu CI, Tang KP, Wang YC, Chiu CH. Impacts of early clinical exposure on undergraduate student professionalism-a qualitative study. *BMC Med Educ*. 2022; 22(1):435. doi:10.1186/s12909-022-03505-5.
16. Ingale MH, Tayade MC, Bhamare S. Early clinical exposure: dynamics, opportunities, and challenges in modern medical education. *J Educ Health Promot*. 2023; 12:295. doi:10.4103/jehp.jehp_237_23.



Ticagrelor versus Clopidogrel in Complex Percutaneous Coronary Intervention in terms of Mortality and Stent Thrombosis

Mujaddid Mudassir, Tanvir Ahmad Raja, Hamza Azhar Ghauri, Muhammad Umar Farooq, Ephraim Sheraz, Muhammad Husnain Yousaf

ABSTRACT

Objective: To compare the effects of ticagrelor and clopidogrel on major adverse cardiovascular events (MACE), 30 day mortality, and stent thrombosis in patients with ST-elevation myocardial infarction (STEMI) undergoing complex percutaneous coronary intervention (PCI).

Methodology: This quasi-experimental study was conducted at the Rawalpindi Institute of Cardiology, Rawalpindi in 3 months from June to August 2025 after ethical approval. After obtaining informed written consent, 376 patients with STEMI treated with complex PCI were included by non-probability convenience sampling technique. The patients were assigned to two groups: Group A received aspirin & clopidogrel, whereas group B received aspirin and ticagrelor. The outcomes assessed were MACE, 30 day mortality, and stent thrombosis. Follow-up was done till 1 month. Data analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 25.

Results: Bifurcation stenting was present in 80(42.6%) patients in clopidogrel group, while it was seen in 46(24.5%) patients in ticagrelor group. The left main PCI was also more prevalent in the ticagrelor group [81(43.1%)]. The thrombolysis in myocardial infarction (TIMI) 3 flow was restored in 53.7% versus 30.3% of the patients taking ticagrelor and clopidogrel, respectively. Stent thrombosis and MACE were significantly higher in the clopidogrel group (8.5% versus 2.7%) and (19.1% versus 11.2%), respectively. In contrast, no substantial difference was noted regarding 30 day mortality (9% vs. 8%) between the two groups.

Conclusion: Ticagrelor is associated with improved post-PCI TIMI flow, lower incidence of MACE and stent thrombosis in STEMI patients who underwent complex PCI as compared to clopidogrel. However, 30 day mortality is not significantly different in the two groups.

Keywords: Ticagrelor. Clopidogrel. Percutaneous coronary intervention.

INTRODUCTION

Primary percutaneous coronary intervention (P-PCI) with insertion of stent is the preferred therapy in cases of myocardial infarction with ST-elevation. However, stent thrombosis (ST) continues to be a rare but catastrophic event that can present in the initial 30 days after implantation.¹ Aspirin and clopidogrel are given as conventional antiplatelet drugs. Aspirin inhibits the enzyme cyclooxygenase-1, thereby stops thromboxane A₂ production, which facilitates platelet plug formation. Clopidogrel is a P₂Y₁₂ receptor inhibitor. These receptors are present on platelets and bind adenosine diphosphate (ADP). So, P₂Y₁₂ inhibitors block the ADP-mediated platelet activation and hence clot formation.^{2,3}

One of the disadvantages of clopidogrel is its variable efficacy. Even with ongoing oral

antiplatelet therapy, atherothrombotic events still occur in some patients. Evidence from several studies suggests that residual platelet activity may be associated with these clinical outcomes, implying that “resistance” to antiplatelet therapy could play a role.⁴ Long-term dual antiplatelet therapy decreases the occurrence of ischemic complications, but is also related to high bleeding risk, contributing to greater morbidity, mortality, and increasing healthcare costs.⁵ The new P₂Y₁₂ receptor antagonist, ticagrelor, has good oral bioavailability. It has a quicker onset of action, more potency, and reversibly inhibits platelet aggregation when compared to conventional P₂Y₁₂ inhibitors like clopidogrel and prasugrel.⁶

This study was conducted to compare the effects of ticagrelor and clopidogrel on major adverse cardiovascular events, 30 day mortality, and stent thrombosis in cases with STEMI undergoing complex PCI. Having more efficacy, ticagrelor would be linked with reduced frequency of complications and mortality. Over the past few years, there have been controversial results from complex PCI cohorts. In some studies, it has been reported that ticagrelor lowers the incidence of MACE than clopidogrel in complex PCI, whereas others claim that the two drugs are not significantly different. These uncertainties underline the

Sharif Medical & Dental College, Sharif Medical City.
Sharif Medical City Road, Off Raiwind Road, Jati Umra,
Lahore 54000, Pakistan.

Correspondence: Dr. Mujaddid Mudassir
Senior Registrar, Department of Cardiology
Rawalpindi Institute of Cardiology, Rawalpindi
E-mail: mujaddidmudassir26@gmail.com

Received: September 27, 2025; Accepted: October 27, 2025

requirement for additional assessment of antiplatelet treatment in this high-risk population.

METHODOLOGY

This quasi-experimental study was carried out at the Rawalpindi Institute of Cardiology, Rawalpindi in 3 months from June to August 2025 after ethical approval from the institution review board (Letter No. RIC/RERC/14/25, 17-07-2025). The sample size of 376 was determined using the incidence of stent thrombosis of 5% with ticagrelor and 13.3% with clopidogrel, 80% power, and 95% confidence interval.⁷ After acquiring written informed consent, STEMI cases treated with complex PCI were included by non-probability convenience sampling technique. Patients of both genders and age >18 years were eligible for inclusion. The diagnosis of STEMI was made from clinical history and ECG findings. Patients complaining of typical chest pain characteristic of myocardial ischemia and a duration of ≥ 20 minutes along with either ST-segment elevation of ≥ 1 mm in two contiguous limb leads, or ≥ 2 mm in two contiguous chest leads, or a new left bundle branch block, were diagnosed with STEMI.⁸ Complex PCI was defined by the presence of any of the following: chronic total occlusion, >60 mm stent length, bifurcation stenting, left main PCI, bypass graft, use of multiwires, atherectomy, guiding catheter extensions or multiple stents.⁹ The exclusion criteria were patients with known allergy or intolerance of antiplatelet therapy, concomitant anticoagulant use, prior bare-metal stent implantation, and history of coronary artery bypass grafting. The patients were assigned to two groups, with 188 patients in each: group A received 75 mg of aspirin once daily and 75 mg of clopidogrel once daily, whereas group B received 75 mg of aspirin once daily and 90 mg of ticagrelor twice daily. Emergency diagnostic coronary angiography and immediate P-PCI were done if the duration from symptom onset was less than 12 hours. The interventional approach (arterial access, stent type, aspiration thrombectomy) was at the discretion of Interventional Cardiologists. The TIMI flow was evaluated in the infarcted vessel pre- and post-procedure. Afterwards, patients were continuously monitored for at least 1 day, and were hospitalized for a minimum of 72 hours. The outcomes assessed were MACE, 30 day mortality, and stent thrombosis. Follow-up was done after 1 month. Major adverse

cardiovascular events included myocardial infarction, stroke, or cardiovascular death.¹⁰

STATISTICAL ANALYSIS

Data were analyzed using the Statistical Package for the Social Sciences (SPSS) version 25. Categorical data was presented as frequencies & percentages, and continuous variables as mean $\pm$ standard deviation (SD). The chi-square test was used for group comparisons and a p-value <0.05 was taken as statistically significant.

RESULTS

The mean age of participants was 58.4 $\pm$ 10.2 years in the clopidogrel group and 57.9 $\pm$ 9.8 years in the ticagrelor group. Patients had a mean body mass index (BMI) of 26.1 $\pm$ 3.7 kg/m² in the clopidogrel group and 26.4 $\pm$ 3.5 kg/m² in the ticagrelor group. The mean ejection fraction was 46.2 $\pm$ 8.5 and 47.1 $\pm$ 8.2 in the clopidogrel and ticagrelor groups, respectively. Regarding gender, the clopidogrel group contained 50.6% of males and 49.4% of females, whereas the ticagrelor group comprised 49.4% of males and 50.6% of females. Diabetes mellitus was more common in the clopidogrel group (50% vs. 39.4%) and so was hypertension (53.3% vs. 46.7%). In the clopidogrel group, 44.4% of the patients had hyperlipidemia versus 47.2% of the patients in the ticagrelor group. The proportion of smokers was 46.1% in the clopidogrel and 47.8% in the ticagrelor group. The history of previous PCI was positive in 50% and 49.4% of the cases in the clopidogrel and ticagrelor groups, respectively.

The procedural characteristics of the two groups are given in Table 1. Significant differences were seen in arterial access, post-PCI TIMI flow 3, bifurcation stenting, and left main PCI.

Major adverse cardiovascular events were frequent in the clopidogrel group [36(19.1%)] as compared to the ticagrelor group [21(11.2%)] with a p-value=0.03. In contrast, no substantial difference was noted in 30 day mortality (9% vs. 8%; p-value=0.71) between the two groups. In terms of the stent thrombosis, the clopidogrel group had a significantly higher number of cases (8.5%) versus the ticagrelor group (2.7%) (p=0.01). These data emphasize that despite a similar incidence of mortality between the two groups, ticagrelor was related with a reduced incidence of MACE and stent thrombosis, suggesting a possible clinical benefit to clopidogrel (Table 2).

Table 1: Procedural Characteristics of the Study Participants

Variables		Clopidogrel Group (n=188)	Ticagrelor Group (n=188)	Total	p-value
		Frequency and Percentage			
Arterial Access	Radial	80(42.6%)	100(53.2%)	180(47.9%)	0.04*
	Femoral	108(57.4%)	88(46.8%)	196(52.1%)	
Severity of Coronary Artery Disease (CAD)	Single Vessel CAD (SVCAD)	68(36.2%)	63(33.5%)	131(34.8%)	0.79
	Double Vessel CAD (DVCAD)	53(28.2%)	52(27.7%)	105(28%)	
	Triple Vessel CAD (TVCAD)	67(35.6%)	73(38.8%)	140(37.2%)	
Treated Vessel	Left Anterior Descending	55(29.3%)	47(25%)	102(27.1%)	0.69
	Left Main	44(23.4%)	44(23.4%)	88(23.5%)	
	Left Circumflex	47(25%)	56(29.8%)	103(27.4%)	
	Right Coronary	42(22.3%)	41(21.8%)	83(22%)	
Type of Stent	Drug Eluting Stents (DES)	95(50.5%)	88(46.8%)	183(48.7%)	0.47
	Bare Metal Stents	93(49.5%)	100(53.2%)	193(51.3%)	
GP2b/3a Inhibitor use	Yes	90(47.9%)	91(48.4%)	181(48.1%)	0.92
	No	98(52.1%)	97(51.6%)	195(51.9%)	
Pre-PCI TIMI Flow	0	51(27.1%)	41(21.8%)	92(24.5%)	0.23
	1-3	137(72.9%)	147(78.2%)	284(75.5%)	
Post-PCI TIMI Flow 3	Yes	57(30.3%)	101(53.7%)	158(42%)	<0.00001*
	No	131(69.7%)	87(46.3%)	218(58%)	
Bifurcation Stenting	Yes	46(24.5%)	80(42.6%)	126(33.5%)	<0.0002*
	No	142(75.5%)	108(57.4%)	250(66.5%)	
Left Main PCI	Yes	45(23.9%)	81(43.1%)	126(33.5%)	0.00008*
	No	143(76.1%)	107(56.9%)	250(66.5%)	
Bypass Graft as Target	Yes	98(52.1%)	83(44.1%)	181(48.1%)	0.12
	No	90(47.9%)	105(55.9%)	195(51.9%)	

*Significant p-value

Table 2: Clinical Outcomes in Patients Taking Clopidogrel versus Ticagrelor

Outcomes		Clopidogrel Group (n=188)	Ticagrelor Group (n=188)	Total	p-value
		Frequency and Percentage			
MACE	Yes	36(19.1%)	21(11.2%)	57(15.2%)	0.03*
	No	152(80.9%)	167(88.8%)	319(84.8%)	
30 day Mortality	Yes	17(9%)	15(8%)	32(8.5)	0.71
	No	171(91%)	173(92%)	344(91.5%)	
Stent Thrombosis	Yes	16(8.5%)	5(2.7%)	21(5.6%)	0.01*
	No	172(91.5%)	183(97.3%)	355(94.4%)	

*Significant p-value

DISCUSSION

Patients with acute coronary syndrome receive conventional antiplatelet regimen consisting of aspirin and P2Y12 receptor inhibitor. However, determining the optimal agent among available P2Y12 inhibitors remains a subject of debate.¹¹ In our study, patients had an average age of 58.2±10 years and the average BMI of 26.3±3.6 kg/m². The males constituted 50% of the study population. There were 44.7% diabetics, 50% hypertensive, 45.8% dyslipidemic, 46.9% smokers, and 49.7% had

previous PCI. In a study, patients had a mean age of 59.1±10.1 years with the majority (80.3%) of males. Their mean BMI was 26.1±3.3 kg/m². Most of the patients had dyslipidemia (82.8%) followed by hypertension (63.1%), smoking (59.9%), diabetes mellitus (34.3%), and prior PCI (18.8%).¹² In another study, the average age was 54.8±10.1 years with 87.3% males and the mean BMI was 26.9±12.3 kg/m². There were 38.4% hypertensive, 30.7% diabetic, 39.5% smokers, 0.4% hyperlipidemic, and 9.6% patients with the history of previous PCI.¹³

In our study, femoral access was predominantly used (52.1%), with the majority of the patients having TVCAD (37.2%), followed by SVCAD (34.8%). In a study by Hakeem et al., radial access was used in 89.9% of the patients. Most of the patients had SVCAD (36.2%) and DVCAD (36.5%).¹³ According to our study, 33.5% of the patients had bifurcation lesions and DES were used in 48.7% of the participants. Around 17.7% had bifurcation lesion and DES were used in 94.1% of the study population according to Hakeem et al.¹³ In our study, bifurcation stenting and left main PCI were more frequently performed in patients taking ticagrelor. Another study reported that ticagrelor did not reduce periprocedural adverse events compared to clopidogrel. Bifurcation stenting was done in 64.6% and 43.4% of the cases taking ticagrelor and clopidogrel, respectively, whereas left main stenting was done 32.4% and 49.4% of the cases taking ticagrelor and clopidogrel, respectively.⁹ Xi et al. also found that ticagrelor was associated with complex procedural outcomes.¹²

Our results revealed that pre-PCI TIMI flow showed no significant variation between the groups, whereas TIMI flow after the procedure was significantly improved in patients taking ticagrelor. A trial reported no variation in TIMI flow before and after procedure in the ticagrelor and clopidogrel groups.¹⁴ In our study, there was a significantly marked reduction in MACE (11.2% versus 19.1%) and stent thrombosis (2.7% versus 8.5%) in the patients taking ticagrelor. In contrast, 30 day mortality was almost the same in the two groups (8% versus 9%). Our results were comparable to other studies. The results of a study showed that ticagrelor was linked to lower incidence of all-cause deaths (1.38% versus 2.85%) and MACE (6.53% versus 9.48%) and the findings were statistically significant.¹⁵ In another study, MACE occurred in 8.6% of patients taking ticagrelor in contrast to 11.2% of the patients taking clopidogrel with a significant difference. Cardiac deaths occurred in 1.2% versus 1.7% in the ticagrelor and clopidogrel groups, respectively with statistically insignificant results.¹² Yan et al. concluded a significant reduction in all-cause deaths and stent thrombosis with ticagrelor, but no significant difference was reported in cardiovascular deaths between patients taking ticagrelor and clopidogrel.¹⁶ Qiu et al. and Bari et al. showed a marked reduction in MACE with ticagrelor as compared to clopidogrel.^{17,18} On contrary, another study revealed no statistical variation in MACE (15% versus 11.8%) and deaths (0.2% versus 0%) in patients treated with ticagrelor and clopidogrel,

respectively.⁹ Another study also showed that 30 day all-cause mortality (14.6% versus 13.5%) and MACE (16.1% versus 14.7%) did not differ statistically in patients taking ticagrelor and clopidogrel.¹⁹ Other studies also showed that MACE was not different in the two groups.^{14,20} In a study, the frequency of MACE was 2.2% versus 2.9%, all-cause deaths were 1.5% versus 1.8%, and stent thrombosis was 1.6% versus 2.1% in the ticagrelor and clopidogrel groups, respectively but with no significance.¹³

CONCLUSION

Ticagrelor is associated with improved post-PCI TIMI flow, lower incidence of MACE, and stent thrombosis in STEMI patients who underwent complex PCI as compared to clopidogrel. However, 30 day mortality is not significantly different in the two groups.

LIMITATIONS & RECOMMENDATIONS

Our findings suggest that ticagrelor may be preferable to clopidogrel, particularly in patients undergoing complex PCI. However, further multicenter randomized studies with larger cohorts and longer follow-up durations are needed to verify these results and to more accurately assess mortality, bleeding, and long-term ischemic outcomes. The quasi-experimental design of this study may introduce selection bias. Additionally, bleeding events, which are important when evaluating the safety profile of ticagrelor, were not comprehensively assessed.

Cost-effectiveness analyses are also needed, especially in poor and developing countries, where issues of affordability and drug accessibility are significant. Incorporation of individualized risk assessment tools (ischemic vs. bleeding risk stratification) may help guide the selection and duration of antiplatelet therapy in clinical practice.

Conflict of interest: None.

Source of funding: None.

Authors' Contributions:

M.M: Conceived and supervised the study.

T.A.R: Reviewed study design and clinical interpretation.

H.A.G: Collected data and performed analysis.

M.U.F: Assisted in data collection and literature review.

E.S: Contributed to data analysis and interpretation.

M.H.Y: Edited and finalized the manuscript.

REFERENCES

1. Kumar M, Shar GS, Kumar R, Shaikh AH, Ishaque H, Khan N, et al. The outcome of primary percutaneous coronary intervention in patients with stent thrombosis. *Indian Heart J.* 2022; 74(6):464-8. doi:10.1016/j.ihj.2022.11.003.

2. Verdoia M, Camaro C, Kedhi E, Marcolongo M, Suryapranata H, De Luca G. Dual antiplatelet therapy duration in acute coronary syndrome patients: the state of the art and open issues. *Cardiovasc Ther.* 2020; 2020:6495036. doi:10.1155/2020/6495036.
3. Virk HUH, Escobar J, Rodriguez M, Bates ER, Khalid U, Jneid H, et al. Dual antiplatelet therapy: a concise review for clinicians. *Life (Basel).* 2023; 13(7):1580. doi:10.3390/life13071580.
4. Karabulut D, Kaya A, Abaci O, Karabulut U, Caglar NT, Arat-Ozkan A. The frequency of aspirin and clopidogrel resistance and related factors in patients undergoing elective percutaneous coronary intervention. *E J Cardiovasc Med.* 2021; 9(2):94-104. doi:10.32596/ejcm.galenos.2021-02-012.
5. Alexopoulos D, Varlamos C, Benetou DR. Type and duration of dual antiplatelet therapy in complex percutaneous coronary intervention. *US Cardiol.* 2020; 14:e14. doi:10.15420/usc.2020.13.
6. Wei P, Wang X, Fu Q, Cao B. Progress in the clinical effects and adverse reactions of ticagrelor. *Thrombosis J.* 2024; 22:8. doi:10.1186/s12959-023-00559-3.
7. Rauf MA, Ziyad M, Mahmood N, Ullah R, Ahmad F, Ullah I, et al. Evaluating the safety and efficacy of ticagrelor vs. clopidogrel following percutaneous coronary intervention in chronic coronary disease. *Cureus.* 2025; 17(1):e77610. doi:10.7759/cureus.77610.
8. Moak JH, Muck AE, Brady WJ. ST-segment elevation myocardial infarction mimics: The differential diagnosis of non-acute coronary syndrome causes of ST-segment/T-wave abnormalities in the chest pain patient. *Turk J Emerg Med.* 2024; 24(4):206-17. doi:10.4103/tjem.tjem_137_24.
9. Lattuca B, Mazeau C, Cayla G, Ducrocq G, Guedeney P, Laredo M, et al. Ticagrelor vs clopidogrel for complex percutaneous coronary intervention in chronic coronary syndrome. *JACC Cardiovasc Interv.* 2024; 17(3):359-70. doi:10.1016/j.jcin.2023.12.011.
10. Bosco E, Hsueh L, McConeghy KW, Gravenstein S, Saade E. Major adverse cardiovascular event definitions used in observational analysis of administrative databases: a systematic review. *BMC Med Res Methodol.* 2021; 21(1):241. doi:10.1186/s12874-021-01440-5.
11. Klein LW. Ticagrelor versus clopidogrel for acute coronary syndrome: have things changed? *Am J Cardiol.* 2024; 210:315-6. doi:10.1016/j.amjcard.2023.10.078.
12. Xi Z, Li J, Qiu H, Guo T, Wang Y, Li Y, et al. Ticagrelor vs. clopidogrel after complex percutaneous coronary intervention in patients with stable coronary artery disease. *Front Cardiovasc Med.* 2021; 8:768190. doi:10.3389/fcvm.2021.768190.
13. Hakeem A, Shah JA, Kumar R, Ali A, Lakho AA, Zeeshan H, et al. Twice-daily clopidogrel vs ticagrelor to reduce short-term major adverse cardiovascular events after primary percutaneous coronary intervention: the TADCLOT trial. *J Am Coll Cardiol.* 2025:S0735-1097(25)07549-7. doi:10.1016/j.jacc.2025.08.041.
14. Hamilos M, Kanakakis J, Anastasiou I, Karvounis C, Vasilikos V, Goudevenos J, et al. Ticagrelor versus clopidogrel in patients with STEMI treated with thrombolysis: the MIRTOS trial. *EuroIntervention.* 2021; 16(14):1163-9. doi:10.4244/EIJ-D-20-00268.
15. Zheng MY, Zhang YK, Hu ST, Gu TS, Jiang C, Zhang JK. Ticagrelor vs. clopidogrel in ST elevation myocardial infarction patients with hemoglobin >160. *Eur Heart J.* 2024; 45(1):1677. doi:10.1093/eurheartj/ehae666.1677.
16. Yan Y, Xu H, Zhao Y, Lin S, Zheng Y. Ticagrelor versus clopidogrel in patients with left main coronary artery stenting. *Cardiovasc Drugs Ther.* 2024. doi:10.1007/s10557-024-07636-3.
17. Qiu M, Zhi X, Liu H, Liang Z, Wang B, Xu K, et al. Impact of ticagrelor vs. clopidogrel in patients with acute coronary syndrome undergoing left main percutaneous coronary intervention. *Cardiology Discovery.* 2025; 5(1):39-46. doi:10.1097/CD9.000000000000134.
18. Bari AM, Raza M, Ijaz ul Haq, Saleem Q, Sofia, Anika. Outcome of ticagrelor versus clopidogrel in ACS patients undergoing PCI. *IJBR.* 2024; 2(2): 594-9. doi:10.70749/ijbr.v2i02.251.
19. Zhou C, Zhang M, Zhao Z, Li E, Zhao Y, Luo W, et al. Clopidogrel vs. ticagrelor in ST-elevation myocardial infarction complicated by cardiogenic shock undergoing primary PCI : findings from a national, multicenter registry. *Thromb J.* 2025; 23(1):38. doi:10.1186/s12959-025-00721-z.
20. Koshy AN, Giustino G, Sartori S, Hooda A, Feng Y, Snyder C, et al. Ticagrelor or prasugrel versus clopidogrel in patients undergoing percutaneous coronary intervention for chronic coronary syndromes. *EuroIntervention.* 2023; 18(15):1244-53. doi:10.4244/EIJ-D-22-00654.



EDITORIAL

- Maternal Malnutrition: Burden on Maternal and Neonatal Health in Pakistan** 1
Amna Iqbal Butt

ORIGINAL ARTICLES

- Outcomes of Percutaneous Coronary Intervention in Chronic Total Occlusion among Women** 3
Muhammad Waleed Hassan, Asif Manzoor, Azhar-ul-Hassan Qureshi, Muhammad Wajahat Alam Lodhi, Waqas Mohiuddin, Muwahhid Rasheed, Sohail Aziz
- Comparison of Soft Tissue Healing in Immediate Implant Placement in Fresh Extraction Socket with and without the Use of Platelet-Rich Fibrin** 9
Tehmina Maryam, Babar Pasha, Muhammad Afzal, Muhammad Nazir Khan, Aiema Nawaz, Noman Tahir
- Diagnostic Accuracy of Alanine Aminotransferase in Intrahepatic Cholestasis of Pregnancy and its Association with Maternal & Fetal/Neonatal Complications** 14
Aisha Yousaf, Nuzhat Sultana, Robina Kouser, Filza Habib, Aurooj Fatima, Hina Zubair
- Association of Serum Cortisol, Interleukin-6, and Serotonin with Depression** 20
Ayesha Tariq, Farhat Bano, Muhammad Haroon Khattak, Muhammad Azhar Sattar, Nasir Wakeel, Aasma Tariq, Shabbir Hussain
- Hepatoprotective Role of Silybum Marianum against Azithromycin-Induced Histological and Biochemical Changes in Albino Wistar Rats** 26
Saba Akram, Mubashra Tahseen, Uzma Bukhari, S. Inayat Ali, Tayyaba Kazmi, Muhammad Mohtasheem ul Hasan
- Cardiac Defects in Neonates Born to Mothers with Gestational Diabetes Mellitus and their Association with Maternal Treatment Regime** 33
Amna Siddique, Muhammad Umar Rasool, Riffat Naeem, Arooj Mirza, Salwa Naeem, Shahid Hamid
- Attitudes of Medical Faculty Enrolled in Masters in Health Professions Education towards Professionalism in a Private Pakistani University: A Mixed-Methods Study** 39
Zaineb Qamar, M. Nasir Ayub Khan, Hina Zaid, Laraib Qamar, Zartashia Arooj, Lamia Yusuf
- Hypoalbuminemia and its Relation with Acute Respiratory Distress Syndrome and In-Hospital Mortality in Intensive Care Unit Patients** 44
Sana Gul, Mohammad Tahir, Rana Ahsan Javed, Hassan Atique, Mehdi Naqvi, Muaz Mubashir
- Association of Clinical and Biochemical Parameters with Renal Resistive Index in Type 2 Diabetic Patients with Normal Albuminuria** 49
Arsalan Shahid, Madiha Naseer, Sudhanshu Kumar Thakur, Mahwish Farzana, Saulat Sarfraz

EDITORIAL

- Rising Rates of Antimicrobial Resistance: A Slow Crisis Demanding Urgent Action** 60
Maria Aslam

ORIGINAL ARTICLES

- Association of HbA1c Levels with Complexity of Coronary Artery Disease in Diabetic Patients Presenting with Acute Coronary Syndrome** 62
Umer Shafiq, Ismail Ahmed Khan, Sohail Aziz, Kashif ur Rehman, Tayyeb Muhammad, Farman Ullah
- Efficacy of Lacosamide in Patients with Diabetic Neuropathy** 68
Awaiz Akram, Shahzad Alam Khan, Umair Ahmad
- Intestinal Parasitic Infections among Children and Young Adolescents Visiting Provincial Headquarter Hospital in Gilgit, Pakistan and their Associated Factors** 73
Uzma Ali, Sadaf Nasir, Adnan Qureshi, Aqsa Aslam, Namra Yunus, Zakia Wahid
- Diagnostic Accuracy of Retrograde Urethrogram in Determination of Urethral Stricture taking Cystoscopy as Gold Standard** 79
Rafia Irum, Aneez Qayyum, Hira Bushra, Hamna Mazhar
- Comparison of Effectiveness between Diclofenac Suppositories and Injection Diclofenac in Abdominal Surgeries** 84
Muhammad Abu Usama, Fizza Laraib, Farhana Manzoor, Maham Fatima, Syed Suleman Zafar Jafary, Hafiz Zain-ul-Abdeen
- Association of Radial Artery Diameter with Radial Artery Spasm in Patients Undergoing Transradial Cardiac Catheterization** 90
Ismail Ahmed Khan, Umer Shafiq, Sohail Aziz, Zohair Aziz, Wajahat Lodhi, Waleed Hassan, Asif Manzoor
- Early Clinical Exposure through Modular Curriculum in Undergraduate Medical Students: Student Experiences** 96
Lamia Yusuf, Aqsa Aslam, Maaz Ahmed, Muhammad Fayyaz Atif, Muhammad Hashim Khan, Muhammad Bin Ahsan
- Ticagrelor versus Clopidogrel in Complex Percutaneous Coronary Intervention in terms of Mortality and Stent Thrombosis** 102
Mujaddid Mudassir, Tanvir Ahmad Raja, Hamza Azhar Ghauri, Muhammad Umar Farooq, Ephraim Sheraz, Muhammad Husnain Yousaf

Subject Index

A

Association of Clinical and Biochemical Parameters with Renal Resistive Index in Type 2 Diabetic Patients with Normal Albuminuria

Arsalan Shahid, 2025; 11(01):49-54

Association of HbA1c Levels with Complexity of Coronary Artery Disease in Diabetic Patients Presenting with Acute Coronary Syndrome

Umer Shafiq, 2025; 11(02):62-67

Association of Radial Artery Diameter with Radial Artery Spasm in Patients Undergoing Transradial Cardiac Catheterization

Ismail Ahmed Khan, 2025; 11(02):90-95

Association of Serum Cortisol, Interleukin-6, and Serotonin with Depression

Ayesha Tariq, 2025; 11(01):20-25

Attitudes of Medical Faculty Enrolled in Masters in Health Professions Education towards Professionalism in a Private Pakistani University: A Mixed-Methods Study

Zaineb Qamar, 2025; 11(01):39-43

C

Cardiac Defects in Neonates Born to Mothers with Gestational Diabetes Mellitus and their Association with Maternal Treatment Regime

Amna Siddique, 2025; 11(01):33-38

Comparison of Effectiveness between Diclofenac Suppositories and Injection Diclofenac in Abdominal Surgeries

Muhammad Abu Usama, 2025; 11(02):84-89

Comparison of Soft Tissue Healing in Immediate Implant Placement in Fresh Extraction Socket with and without the Use of Platelet-Rich Fibrin

Tehmina Maryam, 2025; 11(01):9-13

D

Diagnostic Accuracy of Alanine Aminotransferase in Intrahepatic Cholestasis of Pregnancy and its Association with Maternal & Fetal/Neonatal Complications

Aisha Yousaf, 2025; 11(01):14-19

Diagnostic Accuracy of Retrograde Urethrogram in Determination of Urethral Stricture taking Cystoscopy as Gold Standard

Rafia Irum, 2025; 11(02):79-83

E

Early Clinical Exposure through Modular Curriculum in Undergraduate Medical Students: Student Experiences

Lamia Yusuf, 2025; 11(02):96-101

Efficacy of Lacosamide in Patients with Diabetic Neuropathy

Awais Akram, 2025; 11(02):68-72

H

Hepatoprotective Role of Silybum Marianum against Azithromycin-Induced Histological and Biochemical Changes in Albino Wistar Rats

Saba Akram, 2025; 11(01):26-32

Hypoalbuminemia and its Relation with Acute Respiratory Distress Syndrome and In-Hospital Mortality in Intensive Care Unit Patients

Sana Gul, 2025; 11(01):44-48

I

Intestinal Parasitic Infections among Children and Young Adolescents Visiting Provincial Headquarter Hospital in Gilgit, Pakistan and their Associated Factors

Uzma Ali, 2025; 11(02):73-78

M

Maternal Malnutrition: Burden on Maternal and Neonatal Health in Pakistan

Amna Iqbal Butt, 2025; 11(01):1-2

O

Outcomes of Percutaneous Coronary Intervention in Chronic Total Occlusion among Women

Muhammad Waleed Hassan, 2025; 11(01):3-8

R

Rising Rates of Antimicrobial Resistance: A Slow Crisis Demanding Urgent Action

Maria Aslam, 2025; 11(02):60-61

T

Ticagrelor versus Clopidogrel in Complex Percutaneous Coronary Intervention in terms of Mortality and Stent Thrombosis

Mujaddid Mudassir, 2025; 11(02):102-106

Author Index

Aasma Tariq, 20
Adnan Qureshi, 73
Aiema Nawaz, 9
Aisha Yousaf, 14
Amna Iqbal Butt, 1
Amna Siddique, 33
Aneeza Qayyum, 79
Aqsa Aslam, 73, 96
Arooj Mirza, 33
Arsalan Shahid, 49
Asif Manzoor, 3, 90
Aurooj Fatima, 14
Awais Akram, 68
Ayesha Tariq, 20
Azhar-ul-Hassan Qureshi, 3
Babar Pasha, 9
Ephraim Sheraz, 102
Farhana Manzoor, 84
Farhat Bano, 20
Farman Ullah, 62
Filza Habib, 14
Fizza Laraib, 84
Hafiz Zain-ul-Abdeen, 84
Hamna Mazhar, 79
Hamza Azhar Ghauri, 102
Hassan Atique, 44
Hina Zaid, 39
Hina Zubair, 14
Hira Bushra, 79
Ismail Ahmed Khan, 62, 90
Kashif ur Rehman, 62
Lamia Yusuf, 39, 96
Laraib Qamar, 39
M. Nasir Ayub Khan, 39
Maaz Ahmed, 96
Madiha Naseer, 49
Maham Fatima, 84
Mahwish Farzana, 49
Maria Aslam, 60
Mehdi Naqvi, 44
Mohammad Tahir, 44
Muaz Mubashir, 44
Mubashra Tahseen, 26
Muhammad Abu Usama, 84
Muhammad Afzal, 9
Muhammad Azhar Sattar, 20
Muhammad Bin Ahsan, 96
Muhammad Fayyaz Atif, 96
Muhammad Haroon Khattak, 20
Muhammad Hashim Khan, 96
Muhammad Husnain Yousaf, 102
Muhammad Mohtasheem ul Hasan, 26
Muhammad Nazir Khan, 9
Muhammad Umar Farooq, 102
Muhammad Umar Rasool, 33
Muhammad Wajahat Alam Lodhi, 3
Muhammad Waleed Hassan, 3
Mujaddid Mudassir, 102
Muwahhid Rasheed, 3
Namra Yunus, 73
Nasir Wakeel, 20
Noman Tahir, 9
Nuzhat Sultana, 14
Rafia Irum, 79
Rana Ahsan Javed, 44
Riffat Naeem, 33
Robina Kouser, 14
S. Inayat Ali, 26
Saba Akram, 26
Sadaf Nasir, 73
Salwa Naeem, 33
Sana Gul, 44
Saulat Sarfraz, 49
Shabbir Hussain, 20
Shahid Hamid, 33
Shahzad Alam Khan, 68
Sohail Aziz, 3, 62, 90
Sudhanshu Kumar Thakur, 49
Syed Suleman Zafar Jafary, 84
Tanvir Ahmad Raja, 102
Tayyaba Kazmi, 26
Tayyeb Muhammad, 62
Tehmina Maryam, 9
Umair Ahmad, 68
Umer Shafiq, 62, 90
Uzma Ali, 73
Uzma Bukhari, 26
Wajahat Lodhi, 90
Waleed Hassan, 90
Waqas Mohiuddin, 3
Zaineb Qamar, 39
Zakia Wahid, 73
Zartashia Arooj, 39
Zohair Aziz, 90

Instructions to Authors

The Journal of Sharif Medical & Dental College (JSMDC) agrees to accept manuscripts prepared in accordance with the 'Uniform Requirements for a manuscript submitted to the Biomedical Journals' as approved by the International Committee of Medical Journal Editors (ICMJE) guidelines, published in the British Medical Journal 1991; 302:334-41. All manuscripts must be submitted online at <http://www.ojs.jsmdc.pk>.

FREQUENCY OF PUBLICATION:

JSMDC publishes biannually in June and December.

AIMS & SCOPE:

- JSMDC is a peer-reviewed scientific journal, aims to publish the latest research to improve healthcare outcomes.
- It publishes editorials, original research articles, systematic reviews & meta-analysis, review articles, systematic review articles, photo essays, case reports, recent advances, adverse drug reports, current practices, short communications, and audit reports.
- Studies more than three years old at the time of submission are not entertained as per journal policy. Any study ending three years before the date of submission is judged by the Editorial Board for its suitability as many changes take place over the time period, subject to the area of the study.

ETHICAL/LEGAL CONSIDERATIONS:

The research should be prepared under strict observation of research and publication ethics guidelines recommended by the Council of Science Editors, International Committee of Medical Journal Editors, and World Association of Medical Editors.

- A submitted manuscript must be an original contribution not previously published (except as an abstract or preliminary report), must not be under consideration for publication elsewhere, and if accepted, it must not be published elsewhere in a similar form.
- Each person listed as an author is expected to have participated in the study according to ICMJE criteria. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the Journal, its editors, or the publisher.
- Any study including human subjects or human data must be reviewed and approved by the responsible Institutional Review Board (IRB).
- If the author wishes to publish a paper from his/her dissertation, besides synopsis approval from the CPSP/University, IRB approval is also mandatory.
- Animal experiments should also be reviewed by an appropriate committee (IACUC: Institutional Animal Care and Use Committee) for the care and use of animals. JSMDC follows guidelines for research conducted on animals available at: WMA statement on animal use in biomedical research and the International Association of Veterinary Editors' Consensus Author Guidelines on Animal Ethics and Welfare.
- Randomized Controlled Trial (RCT) should be registered and the trial registration number is mandatory.
- Authors must submit written permission from the copyright owner to use direct quotations, tables, or illustrations that have appeared in copyright form elsewhere, along with complete details about the source.

Instructions to Authors

CONFLICT OF INTEREST:

- JSMDc follows the International Committee of Medical Journal Editors guidelines available at www.icmje.org.
- The authors must provide a formal statement including any potential conflict of interest at the time of submitting the article.
- In case of any conflict of interest, the author must submit an ICMJE form for disclosure of potential conflicts of interest available at: <http://www.jsmdc.pk>.

FINANCIAL DISCLOSURE:

- Each author warrants that he or she has no commercial associations (e.g., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might post a conflict of interest in connection with the submitted article, except as disclosed in a separate attachment.
- All funding sources supporting the work and all institutional or corporate affiliations of the authors are acknowledged in a footnote in the work.

PATIENT ANONYMITY AND INFORMED CONSENT:

It is the author's responsibility to ensure that a patient's anonymity is carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated.

PUBLICATION MISCONDUCT:

- Plagiarism and other publication misconduct, fabrication (picture as well), falsification, duplicate submission, redundant publication, multiple submissions, selective and misleading reporting, selective and misleading referencing are liable to strict action.
- All publication misconducts are dealt with first asking for an explanation from the corresponding author and then from all other authors.
- In case of non-response or unsatisfactory response from the authors, the manuscript is **rejected** if unpublished and **retracted** if published.
- Due notice of retraction will be given in print and on the website.
- The authors will be blacklisted for further submissions and considerations at the journal.
- The authors' Institutional Head will also be informed of the misconduct and relevant action in such a case.

ARTICLE PUBLISHING AND JOURNAL SUBSCRIPTION CHARGES:

- JSMDc is an open-access journal and levies an article processing fee of PKR. 5000/- for the original research article/review article/case report/photo essay.
- The journal will charge an article publishing fee of PKR. 10000/- for the original research article upon acceptance of the submitted article.
- A non-refundable bank draft in favor of Sharif Medical & Dental College A/C No. 15800002100303, HBL SECT branch should be sent via TCS to the editor JSMDc.
- Annual subscription charges of JSMDc are PKR. 2000/-.

These Annual subscription charges are applicable when an individual or a library specifically requests printed copies of the journal. This fee is solely meant to cover printing and postage expenses.

Instructions to Authors

AUTHORSHIP CRITERIA:

- Authors should meet all four criteria for authorship as recommended by the International Committee of Medical Journal Editors (ICMJE). The ICMJE recommends that authorship is based on:
 - Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND
 - Drafting the work or revising it critically for important intellectual content; AND
 - Final approval of the version to be published; AND
 - Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.
- Each author must sign a statement attesting that he or she fulfills the authorship criteria of the uniform requirements. Up to six authors are allowed in a single-institution study. In a multi-institution and international collaborative research, the Editorial Board shall guide on an individual case basis.
- JSMDC strongly discourages gift authorship.
- The author's certification proforma should be filled and submitted along with the manuscript.

Corresponding Author: The authors must assign a corresponding author. The corresponding author is responsible for manuscript corrections, revisions, proofreading & resubmission of revised manuscripts till the acceptance of the manuscript.

ACKNOWLEDGEMENT:

- Individuals who participated in the development of a manuscript but do not qualify as an author should be acknowledged.
- Organizations/departments that provided support in terms of funding and/or other resources should also be acknowledged.

PLAGIARISM POLICY:

Publications are the end-products of the scientific work and their quality reflects the research work. A published paper becomes a source for references, post-publication review, and critique. To contribute to the bulk of knowledge, the paper should be credible.

- JSMDC follows the ICMJE and HEC criteria for all types of plagiarism. The criteria can be accessed at www.icmje.org and www.hec.gov.pk.
- All the submitted manuscripts will be checked for potential plagiarism by "TURNITIN" software. Articles with a similarity index of more than 19% and any single source similarity index of $\geq 5\%$ will not be published. The manuscript will be sent back to the author for revision to bring the similarity index down to the permissible limit. After revision, it will again be checked through the "TURNITIN" for plagiarism. If the similarity report is $< 19\%$, it will be processed further. The article will be sent back to the author again for corrections if the similarity index is still more than 19%. If the author fails to rectify after the second revision, then the manuscript is rejected.
- If the submitted article is found to have been totally or substantially plagiarized then the article is rejected outright and the corresponding author will be informed regarding this misconduct. The author will be asked to give a written explanation.
- For published articles, the allegedly plagiarized article will be temporarily retracted from the issue and a notice to the effect will be published in the JSMDC. The author will be asked for an explanation. In case of unsatisfactory explanation or non-response, the article will be permanently retracted and the author will be blacklisted.

Instructions to Authors

PEER-REVIEW POLICY:

- JSMDC follows a double-blind peer-review process in which both the reviewer and the author remain anonymous throughout the process of review.
- Reviewers are selected according to their specialty and expertise.
- The reviewer database is updated regularly.
- Each manuscript is sent to two external peer reviewers.
- Once the reviewed manuscript is received from both the reviewers, their comments/suggestions are communicated to the author for correction.
- The revised manuscript received from the author is re-assessed by the reviewer/editor and the final decision regarding article acceptance/rejection is also made by the editor.

MANUSCRIPT SUBMISSION GUIDELINES:

ORIGINAL ARTICLES: Original articles should report original research of relevance to clinical medicine. The article should not exceed 4500 words in length (excluding title page, abstract, tables, figures, and references). The original article should follow the basic structure of abstract, introduction, methodology, statistical analysis, results, discussion, conclusion, limitations & recommendations of the study, and references.

Abstract: It should contain a structured abstract of about 250 words. The abstract must have four sections: objectives, methodology, results, and conclusion. Three to five keywords should be given as per Medical Subject Heading (MeSH).

Introduction: The introduction should clearly describe the topic with a brief literature review. All abbreviations used should be first preceded by the word for which it stands. At the end of the introduction, mention the rationale or scientific significance of the study.

Methodology: The methodology should include the study design, place of the study conducted, sample size, sampling technique, ethical consideration, consent, and method & equipment used for research work.

Statistical Analysis: Mention the software used for data analysis and the statistical tests applied.

Results: Text, tables, and figures should be used for illustration of results. Figures and tables should be numbered in order of appearance in the text. Legends for the tables and figures should be typed double-spaced and should include a description of the features shown. All numbers and percentages should be triple-checked. Previously published material and figures should include permission to reproduce from the original publication and original author. Photographs with faces should be accompanied by permission to publish from the subject of the photograph or by a parent in case of a minor. Colored photographs should be submitted in JPEG format.

Discussion: The results of the study should be interpreted with reference to other latest studies.

Conclusion: The conclusion should be clear and concise.

Limitations of the study

Recommendations of the study

References: References must be numbered consecutively according to their appearance in the text. References should be cited in the correct “Vancouver style” with a DOI number. List all authors if the total number of authors is six or less and for more than six authors use et al. after six. Journal names should be abbreviated according to the Index Medicus/MEDLINE. The date of access should be provided for online citations. References should not be older than the last five years.

CLINICAL CASE REPORTS: Must be of academic & educational value and provide relevance to the disease being reported as unusual. It should have a non-structured abstract of about 100-150 words (case-specific) with around 6-8 references and 3 keywords.

Instructions to Authors

LETTER TO THE EDITOR (LTE): It is usually a type of short communication that can be written on any topic that attracts the attention of the reader. There are different types of letters to the editor. If the purpose of the LTE is to comment on a published article, the first sentence of the LTE should include the name of the published article's first author along with the title of the published article and then the comments. If the LTE is a reply to a previously submitted LTE, the first sentence should include the name of the letter's author and cite the letter as a reference. The previously published article should then be referenced as well either in the body of the text or at the end of the response to the LTE.

PHOTO ESSAYS: The journal accepts manuscripts for consideration as photo essays. These essays include the visual presentation of material where the prime emphasis is on the images. These images can include colored images, angiograms, optical coherence tomography, histologic sections, x-rays, ultrasounds, and other studies. The images can be an outstanding presentation of classic findings, atypical findings, or new findings. These are not case reports, but rather a visual presentation of material as a teaching tool. The images need to be of the highest quality. The accompanying manuscript should be limited to a total of 300 words. A maximum of 6-10 separate images and 6 references can be included.

REVIEW ARTICLE: This should consist of a critical overview/analysis of some relatively narrow topic providing background and the recent development with reference to the original literature. It should incorporate the author's original work on the same subject. The review article should be 2500 to 3000 words in length. It should have a non-structured abstract of 150 words with a minimum of 3 keywords. An author can write a review article only if he/she has written a minimum of three original research articles.

SYSTEMATIC REVIEW ARTICLE: It should consist of a well-defined research question and should provide a detailed review of a specific topic based on the existing literature. It should include the collection and analysis of data from all the relevant research in support of the research question being asked. The text should be 2500-3000 words. It should have a structured abstract with a minimum of three keywords.

META-ANALYSIS: It should comprise a statistical analysis of the combined results of numerous scientific studies addressing the same research question. Meta-analysis is a quantitative and epidemiological study design that should critically analyze the results of previous scientific research, mostly randomized controlled trials.

OTHER SECTIONS: The journal also accepts manuscripts for other sections such as diagnostic & therapeutic challenges, clinicopathological correlations, surgical techniques, and new instruments. Diagnostic & therapeutic challenges require no abstract and have no limit for figures and references. Surgical techniques and clinicopathological correlations are treated as a full manuscript and require an abstract. All correspondence and new instruments should have a standard title page with a full-length title, running title, and author information. Keywords anti summary statement should be on the second page. An abstract is not required by the journal for correspondence and new instruments. A summary statement of 50 words is necessary for publication and indexing and must be included at the time of submission. All pages must be numbered starting with the title page being page one. Each figure must be submitted separately.



SHARIF TRUST



Sharif Medical & Dental College

Sharif Medical City Road, Off Raiwind Road, Jati Umra, Lahore 54000, Pakistan.

Tel: +92-4237860101-4 Fax: +92-4237860122 UAN- 042-111-123-786

E-mail: editors@jsmdc.pk Web: www.jsmdc.pk